

HARMONIC ANALYSIS OPERATORS ASSOCIATED WITH MULTIDIMENSIONAL BESSEL OPERATORS

J.J. BETANCOR, A.J. CASTRO, AND J. CURBELO

ABSTRACT. In this paper we establish that the maximal operator and the Littlewood-Paley g-function associated with the heat semigroup defined by multidimensional Bessel operators are of weak type (1,1). Also, we prove that Riesz transforms in the multidimensional Bessel setting are of strong type (p,p), for every $1 < p < \infty$, and of weak type (1,1).

1. INTRODUCTION

Harmonic analysis operators (maximal operators, Riesz transforms, Littlewood-Paley g-functions, multipliers, ...) associated with orthogonal expansions in discrete and continuous settings have been studied since the 60's of the last century, but in the last years it has been a very active area and many authors have studied this topic (see, for instance, [1], [10], [11], [12], [13], [15], [16], [20], [21], [22], [25], [26], [27], [28], [32] and [33]). Muckenhoupt and Stein [22] made a deep study about harmonic analysis in the one dimensional ultraspherical and Bessel settings. We are interested in this paper in the L^p -boundedness properties for the maximal operator and Littlewood-Paley g-function associated with the heat semigroup for the multidimensional Bessel operators and the Riesz transforms in the multidimensional Bessel context. As far as we know harmonic analysis operators in the Bessel settings had been studied always in the one dimensional case. After [22], Andersen [2], Kerman ([17] and [18]) and Andersen and Kerman [3] established L^p -weighted inequalities for the Bessel-Riesz transforms. Stempak [31] studied Littlewood-Paley g-functions and Mihlin-Hörmander multipliers in the Bessel setting. Recently, in [4], [5], [6], [7], [8], and [9] the one-dimensional harmonic analysis in the Bessel context has been completed investigating properties of g-functions and Riesz transforms of every order and maximal operators associated with Poisson and heat semigroups.

We consider the n-dimensional Bessel operator $\Delta_{\lambda_1, \dots, \lambda_n}$ defined by

$$\Delta_{\lambda_1, \dots, \lambda_n} = \sum_{j=1}^n \Delta_{\lambda_j}^{(j)},$$

where $\Delta_{\lambda_j}^{(j)} = -x_j^{-2\lambda_j} \frac{\partial}{\partial x_j} \left(x_j^{2\lambda_j} \frac{\partial}{\partial x_j} \right)$, and $\lambda_j > -1/2$, for every $j = 1, \dots, n$, $n \geq 2$. The heat semigroup generated by $\Delta_{\lambda_1, \dots, \lambda_n}$ is represented by $\{W_t^{\lambda_1, \dots, \lambda_n}\}_{t>0}$. This semigroup is a symmetric diffusion semigroup in the sense of [29] with respect to the measure $m_{\lambda_1, \dots, \lambda_n}$ defined by $\prod_{j=1}^n x_j^{2\lambda_j} dx$ on $(0, \infty)^n$. Then, according to [29, p. 73] the maximal operator

$$W_*^{\lambda_1, \dots, \lambda_n}(f) = \sup_{t>0} \left| W_t^{\lambda_1, \dots, \lambda_n}(f) \right|$$

is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$. Inspired in the ideas developed by Nowak and Sjögren in [23] we establish that $W_*^{\lambda_1, \dots, \lambda_n}$ is of weak type (1,1) with respect to the measure $m_{\lambda_1, \dots, \lambda_n}$.

Theorem 1.1. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. The maximal operator $W_*^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

The first author is partially supported by MTM2007/65609. The second author is supported by a grant for Master studies of "la Caixa". The third author is supported by a grant JAE-Predoc of the CSIC (Spain).

Since for every $f \in C_c^\infty((0, \infty)^n)$, the space of C^∞ -functions on $(0, \infty)^n$ that have compact support,

$$\lim_{t \rightarrow 0^+} W_t^{\lambda_1, \dots, \lambda_n}(f)(x) = f(x), \quad x \in (0, \infty)^n,$$

and $C_c^\infty((0, \infty)^n)$ is a dense subspace of $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, for every $1 \leq p < \infty$, standard arguments allow us to deduce from the L^p -boundedness properties of the maximal operator $W_*^{\lambda_1, \dots, \lambda_n}$ the following result.

Corollary 1.1. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$, and $1 \leq p < \infty$. For every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$,*

$$\lim_{t \rightarrow 0^+} W_t^{\lambda_1, \dots, \lambda_n}(f)(x) = f(x), \quad \text{a.e. } x \in (0, \infty)^n.$$

According to [29, p. 67] the semigroup $\{W_t^{\lambda_1, \dots, \lambda_n}\}_{t>0}$ admits an analytic extension to $\Omega = \{t \in \mathbb{C} : |\operatorname{Arg}(t)| < \pi/4\}$. The Littlewood-Paley g-function of the first order associated with $\{W_t^{\lambda_1, \dots, \lambda_n}\}_{t>0}$ is defined by

$$g^{\lambda_1, \dots, \lambda_n}(f)(x) = \left\{ \int_0^\infty \left| t \frac{\partial}{\partial t} W_t^{\lambda_1, \dots, \lambda_n}(f)(x) \right|^2 \frac{dt}{t} \right\}^{1/2}, \quad x \in (0, \infty)^n.$$

By [29, p. 111], $g^{\lambda_1, \dots, \lambda_n}$ defines a bounded operator from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$. We complete this result analyzing the behavior of $g^{\lambda_1, \dots, \lambda_n}$ on $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Theorem 1.2. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. Then, the operator $g^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

The Bessel operator Δ_λ can be factorized as follows

$$\Delta_\lambda = -x^{-2\lambda} \frac{d}{dx} \left(x^{2\lambda} \frac{d}{dx} \right) = \left(\frac{d}{dx} \right)^* \frac{d}{dx},$$

where $(\frac{d}{dx})^*$ denotes the formal adjoint of $\frac{d}{dx}$ in $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. According to [22, Section 16] and [29] we define, for every $i = 1, \dots, n$, the i -th Riesz transform $R_i^{\lambda_1, \dots, \lambda_n}$ associated with $\Delta_{\lambda_1, \dots, \lambda_n}$ by

$$R_i^{\lambda_1, \dots, \lambda_n} f = \frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f, \quad f \in C_c^\infty((0, \infty)^n).$$

Here $\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2}$ represents the negative square root of $\Delta_{\lambda_1, \dots, \lambda_n}$ whose definition will be specified later (see Section 4). L^p -boundedness properties of the Riesz transforms $R_i^{\lambda_1, \dots, \lambda_n}$, $i = 1, \dots, n$, are established in the following.

Theorem 1.3. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$, and $i = 1, \dots, n$. For every $f \in C_c^\infty((0, \infty)^n)$, $\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f$ admits derivative with respect to x_i on almost all $(0, \infty)^n$ and*

$$\frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad \text{a.e. } x \in (0, \infty)^n,$$

where

$$R_i^{\lambda_1, \dots, \lambda_n}(x, y) = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{\partial}{\partial x_i} W_t^{\lambda_1, \dots, \lambda_n}(x, y) \frac{dt}{\sqrt{t}}, \quad x, y \in (0, \infty)^n,$$

and $W_t^{\lambda_1, \dots, \lambda_n}(x, y)$ represents the kernel of the operator $W_t^{\lambda_1, \dots, \lambda_n}$, for every $t > 0$ (see (1) for definitions).

Moreover, the maximal operator $R_{i,*}^{\lambda_1,\dots,\lambda_n}$ defined by

$$R_{i,*}^{\lambda_1,\dots,\lambda_n}(f)(x) = \sup_{\varepsilon>0} \left| \int_{|x-y|>\varepsilon} R_i^{\lambda_1,\dots,\lambda_n}(x,y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|, \quad x \in (0, \infty)^n,$$

is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. The Riesz transform $R_i^{\lambda_1,\dots,\lambda_n}$ can be extended to $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ by

$$R_i^{\lambda_1,\dots,\lambda_n}(f)(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} R_i^{\lambda_1,\dots,\lambda_n}(x,y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad \text{a.e. } x \in (0, \infty)^n,$$

for every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, that is a bounded operator from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

We now recall some definitions and properties that will be useful in the sequel. If J_ν denotes the Bessel function of the first kind and order $\nu > -1$, we have that, when $\lambda > -1/2$,

$$\Delta_{\lambda,x} \left((xz)^{-\lambda+1/2} J_{\lambda-1/2}(xz) \right) = z^2 (xz)^{-\lambda+1/2} J_{\lambda-1/2}(xz), \quad x, z \in (0, \infty).$$

Then, the heat semigroup $\{W_t^\lambda\}_{t>0}$ generated by $\Delta_{\lambda,x}$ is given by

$$W_t^\lambda(f)(x) = \int_0^\infty W_t^\lambda(x,y) f(y) y^{2\lambda} dy,$$

where

$$W_t^\lambda(x,y) = \int_0^\infty e^{-tz^2} (xz)^{-\lambda+1/2} J_{\lambda-1/2}(xz) (yz)^{-\lambda+1/2} J_{\lambda-1/2}(yz) z^{2\lambda} dz, \quad t, x, y \in (0, \infty).$$

Moreover, according to [34, p. 195], we can write

$$W_t^\lambda(x,y) = \frac{(xy)^{-\lambda+1/2}}{2t} I_{\lambda-1/2} \left(\frac{xy}{2t} \right) e^{-(x^2+y^2)/4t}, \quad t, x, y \in (0, \infty),$$

where I_ν represents the modified Bessel function of the first kind and order $\nu > -1$.

The heat semigroup $\{W_t^{\lambda_1,\dots,\lambda_n}\}_{t>0}$ associated with the multidimensional Bessel operator $\Delta_{\lambda_1,\dots,\lambda_n}$ is defined by

$$W_t^{\lambda_1,\dots,\lambda_n}(f)(x) = \int_{(0,\infty)^n} W_t^{\lambda_1,\dots,\lambda_n}(x,y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy,$$

being

$$(1) \quad W_t^{\lambda_1,\dots,\lambda_n}(x,y) = \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j), \quad x = (x_1, \dots, x_n), \quad y = (y_1, \dots, y_n) \in (0, \infty)^n.$$

The following properties of the modified Bessel function I_ν , $\nu > -1$, will be used (see [19] and [34]). I_ν admits the following series representation

$$(2) \quad I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k+\nu}}{k! \Gamma(\nu+k+1)}, \quad z \in (0, \infty).$$

Then,

$$(3) \quad I_\nu(z) \sim \frac{1}{2^\nu \Gamma(\nu+1)} z^\nu, \quad \text{as } z \rightarrow 0^+.$$

Also,

$$(4) \quad I_\nu(z) = \frac{e^z}{\sqrt{2\pi z}} \left(\sum_{k=0}^m (-1)^k [\nu, k] (2z)^{-k} + O\left(\frac{1}{z^{m+1}}\right) \right), \quad m \in \mathbb{N}, \quad z \in (0, \infty),$$

where $[\nu, 0] = 1$ and

$$[\nu, k] = \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \cdots (4\nu^2 - (2k - 1)^2)}{2^{2k} \Gamma(k + 1)}, \quad k = 1, 2, \dots$$

From (2) it is easy to deduce that

$$(5) \quad \frac{d}{dz} (z^{-\nu} I_\nu(z)) = z^{-\nu} I_{\nu+1}(z), \quad z \in (0, \infty).$$

Asymptotic expansion (4) allows us to connect the operators (maximal operators, g-functions and Riesz transforms) associated with $\Delta_{\lambda_1, \dots, \lambda_n}$ with the corresponding operators in the classical Euclidean setting in a local region that is close to the diagonal in $(0, \infty)^n$. This is a crucial point in the proof of our theorems. The proof of Theorems 1.2 and 1.3 is more involved than the one of Theorem 1.1 because the maximal operator $W_*^{\lambda_1, \dots, \lambda_n}$ is positive while Littlewood-Paley g-function and Riesz transforms are not positive operators.

This paper is organized as follows. In Section 2 we prove Theorem 1.1, Theorem 1.2 is showed in Section 3 and a proof of Theorem 1.3 is presented in Section 4.

Throughout this paper we will use repeatedly without saying it that, for every $k \in \mathbb{N}$, there exists $C_k > 0$ such that $z^k e^{-z} \leq C_k$, $z > 0$. Always by C we denote a suitable positive constant that can change from one line to the other one.

2. PROOF OF THEOREM 1.1

In this section we prove that the maximal operator

$$W_*^{\lambda_1, \dots, \lambda_n}(f)(x) = \sup_{t>0} \left| \int_{(0, \infty)^n} \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|, \quad x \in (0, \infty)^n,$$

where $\lambda_j > -1/2$, $j = 1, \dots, n$, is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

By W_t we denote the classical heat kernel in one dimension, that is,

$$W_t(x, y) = \frac{1}{\sqrt{2\pi t}} e^{-(x-y)^2/4t}, \quad t \in (0, \infty), \quad x, y \in \mathbb{R}.$$

We will split the region $(0, \infty)^n$ of integration in several parts and study each of the operators that appear associated with every part.

Firstly we analyze some auxiliary operators that will be useful in the sequel. The Hardy type operator H_∞ defined by

$$H_\infty(g)(x) = \int_x^\infty \frac{g(y)}{y} dy, \quad x \in (0, \infty),$$

is bounded from $L^1((0, \infty), x^{2\alpha} dx)$ into itself when $\alpha > -1/2$. Then, for every $k \in \mathbb{N}$, the operator

$$H_\infty^k(g)(x) = \int_{x_1}^\infty \cdots \int_{x_k}^\infty \frac{g(y_1, y_2, \dots, y_k)}{y_1 y_2 \cdots y_k} dy_k \cdots dy_1, \quad x = (x_1, \dots, x_k) \in (0, \infty)^k,$$

is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself provided that $\alpha_j > -1/2$, $j = 1, \dots, k$.

Suppose that $k \in \mathbb{N}$ and $\alpha_j > -1/2$, $j = 1, \dots, k$. We define the operator $L_{\alpha_1, \dots, \alpha_k}$ by

$$L_{\alpha_1, \dots, \alpha_k}(g)(x) = \frac{1}{\left(\sum_{j=1}^k x_j\right)^{2 \sum_{j=1}^k (\alpha_j + 1/2)}} \int_0^{x_1} \cdots \int_0^{x_k} g(y) \prod_{j=1}^k y_j^{2\alpha_j} dy, \quad x \in (0, \infty)^k.$$

$L_{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$. Indeed, let $g \in L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ and $\gamma > 0$. By denoting $m_{\alpha_1, \dots, \alpha_k}$ the measure $\prod_{j=1}^k x_j^{2\alpha_j} dx$ we have that

$$\begin{aligned} & m_{\alpha_1, \dots, \alpha_k}(\{x \in (0, \infty)^k : |L_{\alpha_1, \dots, \alpha_k}(g)(x)| > \gamma\}) \\ & \leq m_{\alpha_1, \dots, \alpha_k}\left(\left\{x \in (0, \infty)^k : \left(\sum_{j=1}^k x_j\right)^{-2 \sum_{j=1}^k (\alpha_j + 1/2)} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)} > \gamma\right\}\right) \\ & = m_{\alpha_1, \dots, \alpha_k}\left(\left\{x \in (0, \infty)^k : \sum_{j=1}^k x_j < \left(\frac{1}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)}\right)^{1/(2 \sum_{j=1}^k (\alpha_j + 1/2))}\right\}\right) \\ & \leq m_{\alpha_1, \dots, \alpha_k}(Q), \end{aligned}$$

where $Q = \prod_{j=1}^k [0, (\frac{1}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)})^{1/(2 \sum_{j=1}^k (\alpha_j + 1/2))}]$.

Since,

$$m_{\alpha_1, \dots, \alpha_k}(Q) \leq \frac{C}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)},$$

it concludes that

$$m_{\alpha_1, \dots, \alpha_k}(\{x \in (0, \infty)^k : |L_{\alpha_1, \dots, \alpha_k}(g)(x)| > \gamma\}) \leq \frac{C}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)}.$$

Let $\alpha_j \in \mathbb{R}$, $j = 1, \dots, k$, with $k \in \mathbb{N}$, we define the local Hardy type operator $H_{loc}^{\alpha_1, \dots, \alpha_k}$ by

$$H_{loc}^{\alpha_1, \dots, \alpha_k}(g)(x) = \frac{1}{\prod_{j=1}^k x_j^{2\alpha_j + 1}} \int_{x_1/2}^{2x_1} \cdots \int_{x_k/2}^{2x_k} g(y) \prod_{j=1}^k y_j^{2\alpha_j} dy, \quad x = (x_1, \dots, x_k) \in (0, \infty)^k.$$

It is not hard to see that $H_{loc}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself.

Also we consider the local maximal operator $\mathbb{W}_{*,loc}^{\alpha_1, \dots, \alpha_k}$ defined by

$$\mathbb{W}_{*,loc}^{\alpha_1, \dots, \alpha_k}(g)(x) = \sup_{t>0} \left| \int_{x_1/2}^{2x_1} \cdots \int_{x_k/2}^{2x_k} \prod_{j=1}^k (x_j y_j)^{-\alpha_j} \mathbb{W}_t(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\alpha_j} dy \right|, \quad x \in (0, \infty)^k.$$

$\mathbb{W}_{*,loc}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$. Indeed, for every $j = (j_1, \dots, j_k) \in \mathbb{Z}^k$, we denote by Q_j the dyadic cube $Q_j = \prod_{i=1}^k [2^{j_i}, 2^{j_i+1}]$. Also, for every $x \in (0, \infty)^k$, $D(x)$ represents the following set

$$D(x) = \{y = (y_1, \dots, y_k) \in (0, \infty)^k : x_j/2 < y_j < 2x_j, \quad j = 1, \dots, k\}.$$

Note that, if $x \in Q_j$ and $y \in D(x)$, then $2^{j_i-1} \leq y_i \leq 2^{j_i+2}$, $i = 1, \dots, k$.

Hence, it has

$$\begin{aligned} \mathbb{W}_{*,loc}^{\alpha_1,\dots,\alpha_k}(g)(x) &\leq \sup_{t>0} \int_{2^{j_1-1}}^{2^{j_1+2}} \cdots \int_{2^{j_k-1}}^{2^{j_k+2}} \prod_{i=1}^k (x_i y_i)^{-\alpha_i} \mathbb{W}_t(x_i, y_i) |g(y)| \prod_{i=1}^k y_i^{2\alpha_i} dy \\ &\leq C \sup_{t>0} \int_{2^{j_1-1}}^{2^{j_1+2}} \cdots \int_{2^{j_k-1}}^{2^{j_k+2}} \prod_{i=1}^k \mathbb{W}_t(x_i, y_i) |g(y)| dy, \quad x \in Q_j, \end{aligned}$$

where $C > 0$ does not depend on $j \in \mathbb{Z}^k$.

If we denote, for every $j = (j_1, \dots, j_k) \in \mathbb{Z}^k$, by D_j the set

$$D_j = \{y = (y_1, \dots, y_k) \in (0, \infty)^k : 2^{j_i-1} < y_i < 2^{j_i+2}, \quad i = 1, \dots, k\},$$

since, as it is well known, the maximal operator

$$\mathbb{W}_*(f)(x) = \sup_{t>0} \left| \int_{\mathbb{R}^k} \prod_{i=1}^k \mathbb{W}_t(x_i, y_i) f(y) dy \right|, \quad x \in \mathbb{R}^k,$$

is bounded from $L^1(\mathbb{R}^k, dx)$ into $L^{1,\infty}(\mathbb{R}^k, dx)$, it follows, for every $\gamma > 0$, that

$$\begin{aligned} m_{\alpha_1,\dots,\alpha_k}(\{x \in (0, \infty)^k : \mathbb{W}_{*,loc}^{\alpha_1,\dots,\alpha_k}(g)(x) > \gamma\}) &= \sum_{j \in \mathbb{Z}^k} m_{\alpha_1,\dots,\alpha_k}(\{x \in Q_j : \mathbb{W}_{*,loc}^{\alpha_1,\dots,\alpha_k}(g)(x) > \gamma\}) \\ &\leq C \sum_{j \in \mathbb{Z}^k} 2^{\sum_{i=1}^k 2^{j_i} \alpha_i} m_0^{(k)}(\{x \in Q_j : \mathbb{W}_{*,loc}^{\alpha_1,\dots,\alpha_k}(g)(x) > \gamma\}) \\ &\leq C \sum_{j \in \mathbb{Z}^k} 2^{\sum_{i=1}^k 2^{j_i} \alpha_i} m_0^{(k)}(\{x \in \mathbb{R}^k : \mathbb{W}_*(\chi_{D_j} |g|)(x) > \gamma M\}) \\ &\leq \frac{C}{\gamma} \sum_{j \in \mathbb{Z}^k} 2^{\sum_{i=1}^k 2^{j_i} \alpha_i} \int_{D_j} |g(y)| dy \\ &\leq \frac{C}{\gamma} \sum_{j \in \mathbb{Z}^k} \int_{D_j} |g(y)| \prod_{i=1}^k y_i^{2\alpha_i} dy \\ &\leq \frac{C}{\gamma} \|g(y)\|_{L^1((0,\infty)^k, \prod_{i=1}^k x_i^{2\alpha_i} dx)}, \quad g \in L^1((0, \infty)^k, \prod_{i=1}^k x_i^{2\alpha_i} dx). \end{aligned}$$

Here $m_0^{(k)}$ denotes the usual Lebesgue measure on $\mathbb{R}^k$ and M is a suitable positive constant that does not depend on $j \in \mathbb{Z}^k$.

We now define, for every $1 \leq l \leq k$, $l, k \in \mathbb{N}$, and $\alpha_j > -1/2$, $j = 1, \dots, k$, the operator

$$\begin{aligned} \mathcal{H}_{l,k}^{\alpha_1,\dots,\alpha_k}(g)(x) &= \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_k/2}^{2x_k} \prod_{j=l+1}^k (x_j y_j)^{-\alpha_j} \\ &\quad \times \frac{g(y)}{\left(\sum_{j=1}^k (x_j - y_j)^2 \right)^\varepsilon} \prod_{j=1}^k y_j^{2\alpha_j} dy, \quad x = (x_1, \dots, x_k) \in (0, \infty)^k, \end{aligned}$$

where $\varepsilon = \sum_{j=1}^l (\alpha_j + 1/2) + (k-l)/2$. By proceeding as in [23, Case 3], we can prove that $\mathcal{H}_{l,k}^{\alpha_1,\dots,\alpha_k}$

is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$.

In order to study the operator $W_*^{\lambda_1,\dots,\lambda_n}$ we use the estimates (3) and (4) and divide, for every $j = 1, \dots, n$, the region $(0, \infty) \times (0, \infty)$ in three parts: one of them is close to the diagonal

$\{(x_j, y_j) : x_j = y_j\}$ and the other two ones are far from the diagonal.

Our result will be proved when we see that, for every $0 \leq l \leq m \leq n$, $l, m, n \in \mathbb{N}$ the operator $S_{l,m}^{\lambda_1, \dots, \lambda_n}$ defined by

$$S_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) = \sup_{t>0} \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \int_{2x_{m+1}}^\infty \cdots \int_{2x_n}^\infty \\ \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$.

According to [8, Lemma 4], we have that

$$(6) \quad 0 \leq W_t^\lambda(x, y) \leq C y^{-2\lambda-1}, \quad 2x < y < \infty \text{ and } t > 0,$$

provided that $\lambda > -1/2$. Then, for every $0 \leq l \leq m \leq n$, $l, m, n \in \mathbb{N}$,

$$S_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) \leq C \sup_{t>0} \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \prod_{j=1}^m W_t^{\lambda_j}(x_j, y_j) \\ \times \left(\int_{2x_{m+1}}^\infty \cdots \int_{2x_n}^\infty \frac{|f(y)|}{y_{m+1} \cdots y_n} dy_n \cdots y_{m+1} \right) \prod_{j=1}^m y_j^{2\lambda_j} dy_m \cdots dy_1, \quad x \in (0, \infty)^n.$$

Hence, since the operator H_∞^{n-m} is bounded from $L^1((0, \infty)^{n-m}, \prod_{j=1}^{n-m} x_j^{2\alpha_j} dx)$ into itself, for every $0 \leq m < n$, $m \in \mathbb{N}$, by [14, Proposition 1], we must show that, for every $0 \leq l \leq m \leq n$, $l, m \in \mathbb{N}$, and $\alpha_j > -1/2$, $j = 1, \dots, n$, the operator $\mathcal{S}_{l,m}^{\alpha_1, \dots, \alpha_m}$ defined by

$$\mathcal{S}_{l,m}^{\alpha_1, \dots, \alpha_m}(g)(x) = \sup_{t>0} \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \prod_{j=1}^m W_t^{\alpha_j}(x_j, y_j) |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy, \quad x \in (0, \infty)^m,$$

is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$.

Let $l, m \in \mathbb{N}$ such that $0 \leq l \leq m \leq n$ and let $\alpha_j > -1/2$, $j = 1, \dots, m$. According to (4) we get

$$(7) \quad 0 \leq W_t^\lambda(x, y) \leq C \frac{(xy)^{-\lambda}}{\sqrt{t}} e^{-\frac{(x-y)^2}{4t}} \leq C \frac{(xy)^{-\lambda}}{\sqrt{t}} e^{-\frac{x^2}{16t}} \leq C \frac{(xy)^{-\lambda-1/2}}{\sqrt{t}} x e^{-\frac{x^2}{16t}} \leq C \frac{e^{-\frac{x^2}{20t}}}{t^{\lambda+1/2}},$$

provided that $0 < y < x/2 < \infty$, $t > 0$, $xy/t \geq 1$ and $\lambda > -1/2$.

Also by (3) it obtains

$$(8) \quad 0 \leq W_t^\lambda(x, y) \leq C \frac{1}{t^{\lambda+1/2}} e^{-(x^2+y^2)/4t} \leq C \frac{1}{t^{\lambda+1/2}} e^{-x^2/4t},$$

when $t, x, y \in (0, \infty)$, $xy/t \leq 1$ and $\lambda > -1/2$.

By combining (7) and (8) it follows that, for every $\lambda > -1/2$,

$$(9) \quad 0 \leq W_t^\lambda(x, y) \leq C \frac{1}{t^{\lambda+1/2}} e^{-x^2/20t}, \quad 0 < y < x/2 \text{ and } t > 0.$$

Then,

$$\begin{aligned}
\mathcal{S}_{m,m}^{\alpha_1,\dots,\alpha_m}(g)(x) &\leq C \int_0^{x_1/2} \cdots \int_0^{x_m/2} \sup_{t>0} \frac{e^{-\sum_{j=1}^m x_j^2/20t}}{\sum_{j=1}^m (\alpha_j+1/2)} |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \\
&\leq C \frac{1}{\left(\sum_{j=1}^m x_j^2\right)^{\sum_{j=1}^m (\alpha_j+1/2)}} \int_0^{x_1/2} \cdots \int_0^{x_m/2} |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \\
&\leq CL_{\alpha_1,\dots,\alpha_m}(|g|)(x), \quad x \in (0, \infty)^m.
\end{aligned}$$

Hence, $\mathcal{S}_{m,m}^{\alpha_1,\dots,\alpha_m}$ is a bounded operator from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$.

We now analyze the operator $\mathcal{S}_{0,m}^{\alpha_1,\dots,\alpha_m}$. Again, according to (3) and (4), it gets, if $\lambda > -1/2$,

$$(10) \quad 0 \leq W_t^\lambda(x, y) \leq C \frac{(xy)^{-\lambda}}{\sqrt{t}} e^{-(x-y)^2/4t}, \quad t, x, y \in (0, \infty) \text{ and } \frac{xy}{t} \geq 1,$$

and

$$(11) \quad 0 \leq W_t^\lambda(x, y) \leq C \frac{e^{-(x^2+y^2)/4t}}{t^{\lambda+1/2}} \leq C \frac{1}{x^{2\lambda+1}}, \quad t, x, y \in (0, \infty) \text{ and } \frac{xy}{t} \leq 1.$$

From (10) and (11) it infers that, if $\lambda > -1/2$,

$$(12) \quad 0 \leq W_t^\lambda(x, y) \leq C \left(\frac{(xy)^{-\lambda}}{\sqrt{t}} e^{-(x-y)^2/4t} + \frac{1}{x^{2\lambda+1}} \right), \quad t, x, y \in (0, \infty).$$

Then (12) implies that the operator $\mathcal{S}_{0,m}^{\alpha_1,\dots,\alpha_m}$ is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ when this property is established for the operator $\mathcal{D}_{l,m}^{\alpha_1,\dots,\alpha_m}$ defined, for every $0 \leq l \leq m$, $l \in \mathbb{N}$, by

$$\begin{aligned}
\mathcal{D}_{l,m}^{\alpha_1,\dots,\alpha_m}(g)(x) &= \sup_{t>0} \int_{x_1/2}^{2x_1} \cdots \int_{x_l/2}^{2x_l} \prod_{j=1}^l \frac{(x_j y_j)^{-\alpha_j}}{\sqrt{t}} e^{-(x_j - y_j)^2/4t} \\
&\quad \times \left(\frac{1}{\prod_{j=l+1}^m x_j^{2\alpha_j+1}} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} |g(y)| \prod_{j=l+1}^m y_j^{2\alpha_j} dy_j \right) \prod_{j=1}^l y_j^{2\alpha_j} dy_j, \quad x \in (0, \infty)^m.
\end{aligned}$$

Since the operator $\mathbb{W}_{*,loc}^{\alpha_1,\dots,\alpha_l}$, $0 < l \leq m$, is bounded from $L^1((0, \infty)^l, \prod_{j=1}^l x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^l, \prod_{j=1}^l x_j^{2\alpha_j} dx)$ and the operator $H_{loc}^{\alpha_{l+1},\dots,\alpha_m}$, $0 < l \leq m$, is bounded from $L^1((0, \infty)^{m-l}, \prod_{j=l+1}^m x_j^{2\alpha_j} dx)$ into itself, from [14, Proposition 1], it follows that $\mathcal{D}_{l,m}^{\alpha_1,\dots,\alpha_m}$ is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$, for every $0 \leq l \leq m$, $l \in \mathbb{N}$.

Assume that $0 < l < m$. Since the operator $L_{\alpha_1,\dots,\alpha_k}$, $k \in \mathbb{N}$, is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ and the operator $H_{loc}^{\alpha_1,\dots,\alpha_k}$, $k \in \mathbb{N}$, is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself, by taking into account [14, Proposition 1], (9) and (12), in order to prove that the operator $\mathcal{S}_{l,m}^{\alpha_1,\dots,\alpha_m}$ is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into

$L^{1,\infty}((0,\infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$, it is sufficient to see this property for the operator

$$\begin{aligned} T_{l,m}^{\alpha_1,\dots,\alpha_m}(g)(x) &= \sup_{t>0} \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \prod_{j=1}^l \frac{1}{t^{\alpha_j+1/2}} e^{-x_j^2/20t} \\ &\quad \times \prod_{j=l+1}^m \frac{(x_j y_j)^{-\alpha_j}}{\sqrt{t}} e^{-(x_j-y_j)^2/4t} |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy, \quad x \in (0,\infty)^m. \end{aligned}$$

We have that

$$\begin{aligned} T_{l,m}^{\alpha_1,\dots,\alpha_m}(g)(x) &\leq \sup_{t>0} \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \prod_{j=l+1}^m (x_j y_j)^{-\alpha_j} \\ &\quad \times \frac{e^{-(\sum_{j=1}^l x_j^2 + \sum_{j=l+1}^m (x_j-y_j)^2)/20t}}{\sum_{j=1}^l (\alpha_j+1/2) + (m-l)/2} |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \\ &\leq C \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_m/2}^{2x_m} \prod_{j=l+1}^m (x_j y_j)^{-\alpha_j} \\ &\quad \times \frac{1}{\left(\sum_{j=1}^l x_j^2 + \sum_{j=l+1}^m (x_j - y_j)^2 \right)^{\sum_{j=1}^l (\alpha_j+1/2) + (m-l)/2}} |g(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \\ &\leq C \mathcal{H}_{l,m}^{\alpha_1,\dots,\alpha_m}(|g|)(x), \quad x = (x_1, \dots, x_m) \in (0,\infty)^m. \end{aligned}$$

Then, $T_{l,m}^{\alpha_1,\dots,\alpha_m}$ is bounded from $L^1((0,\infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0,\infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$, because $\mathcal{H}_{l,m}^{\alpha_1,\dots,\alpha_m}$ has this property.

Thus the proof of Theorem 1.1 is finished. $\square$

3. PROOF OF THEOREM 1.2

We are going to see that the Littlewood-Paley g-function associated with the heat semigroup $\{W_t^{\lambda_1,\dots,\lambda_n}\}_{t>0}$ and given by

$$g^{\lambda_1,\dots,\lambda_n}(f)(x) = \left\{ \int_0^\infty \left| t \frac{\partial}{\partial t} W_t^{\lambda_1,\dots,\lambda_n}(f)(x) \right|^2 \frac{dt}{t} \right\}^{1/2}, \quad x \in (0,\infty)^n,$$

defines a bounded operator from $L^1((0,\infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0,\infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, when $\lambda_j > -1/2$, $j = 1, \dots, n$.

For each $x \in (0,\infty)^n$, we consider the local operator $g_{loc}^{\lambda_1,\dots,\lambda_n}$ defined by

$$g_{loc}^{\lambda_1,\dots,\lambda_n}(f)(x) = \left\{ \int_0^\infty t \left| \frac{\partial}{\partial t} \int_{x_1/2}^{2x_1} \cdots \int_{x_n/2}^{2x_n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2}.$$

Minkowski's inequality leads us to

$$\begin{aligned} &|g^{\lambda_1,\dots,\lambda_n}(f)(x) - g_{loc}^{\lambda_1,\dots,\lambda_n}(f)(x)| \\ &\leq \left\{ \int_0^\infty t \left| \frac{\partial}{\partial t} \int_{(0,\infty)^n} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j) (x_j y_j)^{-\lambda_j} \right. \right. \right. \end{aligned}$$

$$\left. \times \frac{e^{-(x_j-y_j)^2/4t}}{2\sqrt{\pi t}} \right) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \Big| dt \Big\}^{1/2}, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.$$

Our proof is divided in two steps:

Claim 1. *The operator $g_{loc}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

Claim 2. *The operator $G^{\lambda_1, \dots, \lambda_n}$ defined by*

$$G^{\lambda_1, \dots, \lambda_n}(f)(x) = \left\{ \int_0^\infty t \left| \frac{\partial}{\partial t} \int_{(0, \infty)^n} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j) (x_j y_j)^{-\lambda_j} \right. \right. \right. \\ \left. \left. \times \frac{e^{-(x_j-y_j)^2/4t}}{2\sqrt{t\pi}} \right) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| dt \Big\}^{1/2}, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

We prove firstly Claim 1. It is sufficient to show that the operator

$$g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x) = \left\{ \int_0^\infty t \left| \int_{x_1/2}^{2x_1} \cdots \int_{x_n/2}^{2x_n} (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{t}} \right) \prod_{j=2}^n (x_j y_j)^{-\lambda_j} \right. \right. \\ \left. \left. \times \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| dt \right\}^{1/2}, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. When the derivative acts on any other factor the operator that appears can be analyzed in a similar way.

For every $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$ the operator $Y_m^{\lambda_1, \dots, \lambda_n}$ is defined by

$$Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) = \chi_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})}(x) \left\{ \int_0^\infty t \left| \int_{(0, \infty)^n} \chi_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})}(y) (x_1 y_1)^{-\lambda_1} \right. \right. \\ \left. \left. \times \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{t}} \right) \prod_{j=2}^n (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| dt \right\}^{1/2} \\ = \chi_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})}(x) \prod_{j=1}^n x_j^{-\lambda_j} \left\{ \int_0^\infty t \left| \int_{(0, \infty)^n} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{t}} \right) \right. \right. \\ \left. \left. \times \prod_{j=2}^n \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{t}} f_m(y) dy \right| dt \right\}^{1/2}, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n,$$

where $f_m(y) = \chi_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})}(y) f(y) \prod_{j=1}^n y_j^{\lambda_j}$, $y = (y_1, \dots, y_n) \in (0, \infty)^n$.

Let $\gamma > 0$. We can write, for each $f \in L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$,

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : g_{1, loc}^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma\}) \\
& \leq m_{\lambda_1, \dots, \lambda_n} \left(\{x \in (0, \infty)^n : |g_{1, loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| \right. \\
& \quad \left. + \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma\} \right) \\
& = m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |g_{1, loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \\
& \quad + m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}) \\
(13) \quad & = m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |g_{1, loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \\
& \quad + \sum_{m \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}).
\end{aligned}$$

For every $m \in \mathbb{Z}^n$, it has

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}) \\
& \leq C 2^{\sum_{j=1}^n m_j \lambda_j} m_0^{(n)}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : g_1(f_m)(x) > M \gamma 2^{\sum_{j=1}^n m_j \lambda_j}\}),
\end{aligned}$$

where $C, M > 0$ are suitable constants that do not depend on $m \in \mathbb{Z}^n$, and g_1 represents the classical Littlewood-Paley g-function defined by

$$g_1(F)(x) = \left\{ \int_0^\infty t \left| \int_{\mathbb{R}^n} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{2\sqrt{\pi t}} \right) \prod_{j=2}^n \frac{e^{-(x_j-y_j)^2/4t}}{2\sqrt{\pi t}} F(y) dy \right|^2 dt \right\}^{1/2}, \quad x \in \mathbb{R}^n.$$

Since, as it is well known, g_1 is a bounded operator from $L^1(\mathbb{R}^n, dx)$ into $L^{1,\infty}(\mathbb{R}^n, dx)$, we deduce that,

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}) \\
& \leq \frac{C}{\gamma} 2^{\sum_{j=1}^n m_j \lambda_j} \|f_m\|_{L^1(\mathbb{R}^n, dx)} \\
& = \frac{C}{\gamma} 2^{\sum_{j=1}^n m_j \lambda_j} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{\lambda_j} dy \\
& \leq \frac{C}{\gamma} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

Hence, we obtain

$$(14) \quad \sum_{m \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : Y_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \frac{\gamma}{2}\}) \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)}.$$

On the other hand, it has that

$$(15) \quad m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \\ = \sum_{m \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : |g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}).$$

Fix $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$. We consider, for every $x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1})$, the set

$$A(x) = \prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2}) \setminus \prod_{j=1}^n [x_j/2, 2x_j).$$

It is clear that

$$A(x) = \bigcup_{l=1}^n (B_l(x) \cup C_l(x)), \quad x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}),$$

where, for every $x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1})$ and $l = 1, \dots, n$,

$$B_l(x) = \{y \in (0, \infty)^n : 2^{m_j-1} \leq y_j \leq 2^{m_j+2}, \quad j = 1, \dots, n, \quad j \neq l; \quad 2^{m_l-1} \leq y_l < x_l/2\}$$

and

$$C_l(x) = \{y \in (0, \infty)^n : 2^{m_j-1} \leq y_j \leq 2^{m_j+2}, \quad j = 1, \dots, n, \quad j \neq l; \quad 2x_l \leq y_l < 2^{m_l+2}\}.$$

We get by using Minkowski's inequality, for every $x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1})$,

$$\begin{aligned} & |g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| \\ & \leq \prod_{j=1}^n x_j^{-\lambda_j} \left\{ \int_0^\infty t \left| \sum_{l=1}^n \left(\int_{B_l(x)} + \int_{C_l(x)} \right) \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1-y_1)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=2}^n \frac{e^{-\frac{(x_j-y_j)^2}{4t}}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \\ & \leq \sum_{l=1}^n \left(\prod_{j=1}^n x_j^{-\lambda_j} \left\{ \int_0^\infty t \left| \int_{B_l(x)} \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1-y_1)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=2}^n \frac{e^{-\frac{(x_j-y_j)^2}{4t}}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \right. \\ & \quad \left. + \prod_{j=1}^n x_j^{-\lambda_j} \left\{ \int_0^\infty t \left| \int_{C_l(x)} \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1-y_1)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=2}^n \frac{e^{-\frac{(x_j-y_j)^2}{4t}}}{\sqrt{t}} f(y) \prod_{j=1}^n y_j^{\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \right) \\ & \leq \sum_{l=1}^n \left(\prod_{j=1}^n x_j^{-\lambda_j} \int_{B_l(x)} |f(y)| \left\{ \int_0^\infty t \left| \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1-y_1)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=2}^n \frac{e^{-\frac{(x_j-y_j)^2}{4t}}}{\sqrt{t}} \right|^2 dt \right\}^{\frac{1}{2}} \prod_{j=1}^n y_j^{\lambda_j} dy \right. \\ & \quad \left. + \prod_{j=1}^n x_j^{-\lambda_j} \int_{C_l(x)} |f(y)| \left\{ \int_0^\infty t \left| \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1-y_1)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=2}^n \frac{e^{-\frac{(x_j-y_j)^2}{4t}}}{\sqrt{t}} \right|^2 dt \right\}^{\frac{1}{2}} \prod_{j=1}^n y_j^{\lambda_j} dy \right). \end{aligned}$$

Since

$$(16) \quad \left| \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{t}} \right) \right| \leq C \frac{e^{-(x_1-y_1)^2/8t}}{t^{3/2}}, \quad t, x_1, y_1 \in (0, \infty),$$

it follows that, for every $x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1})$,

$$(17) \quad |g_{1,loc}^{\lambda_1,\dots,\lambda_n}(f)(x) - Y_m^{\lambda_1,\dots,\lambda_n}(f)(x)| \\ \leq C \sum_{l=1}^n \left(\prod_{j=1}^n x_j^{-\lambda_j} \int_{B_l(x)} |f(y)| \left\{ \int_0^\infty \frac{e^{-\sum_{j=1}^n (x_j-y_j)^2/8t}}{t^{n+1}} dt \right\}^{1/2} \prod_{j=1}^n y_j^{\lambda_j} dy \right. \\ \left. + \prod_{j=1}^n x_j^{-\lambda_j} \int_{C_l(x)} |f(y)| \left\{ \int_0^\infty \frac{e^{-\sum_{j=1}^n (x_j-y_j)^2/8t}}{t^{n+1}} dt \right\}^{1/2} \prod_{j=1}^n y_j^{\lambda_j} dy \right).$$

By taking into account symmetries we only study the first term, $l = 1$, in the last sum. Moreover the analysis of the integrals on $B_1(x)$ and $C_1(x)$ can be made similarly. We estimates only the integral on $B_1(x)$. Let $x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1})$. Since $|x_1 - y_1| = x_1 - y_1 > x_1/2 > 2^{m_1-1}$, when $2^{m_1-1} \leq y_1 < x_1/2$, we have,

$$\begin{aligned} & \prod_{j=1}^n x_j^{-\lambda_j} \int_{B_1(x)} |f(y)| \left\{ \int_0^\infty \frac{e^{-\sum_{j=1}^n (x_j-y_j)^2/4t}}{t^{n+1}} dt \right\}^{1/2} \prod_{j=1}^n y_j^{\lambda_j} dy \\ & \leq C \prod_{j=1}^n x_j^{-\lambda_j} \int_{B_1(x)} \frac{|f(y)|}{(2^{2m_1-2} + \sum_{j=2}^n (x_j - y_j)^2)^{n/2}} \prod_{j=1}^n y_j^{\lambda_j} dy \\ & \leq C \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} \frac{|f(y)|}{(2^{2m_1-2} + \sum_{j=2}^n (x_j - y_j)^2)^{n/2}} dy \\ & = C \int_{\mathbb{R}^{n-1}} \frac{\tilde{f}_m(\bar{y})}{(2^{2m_1-2} + \sum_{j=2}^n (x_j - y_j)^2)^{n/2}} d\bar{y}, \end{aligned}$$

where $\bar{y} = (y_2, \dots, y_n) \in \mathbb{R}^{n-1}$ and $\tilde{f}_m(\bar{y}) = \chi_{\prod_{j=2}^n [2^{m_j-1}, 2^{m_j+2})}(\bar{y}) \int_{2^{m_1-1}}^{2^{m_1+2}} |f(y_1, \bar{y})| dy_1$. Then, by using standard argument it gets

$$\begin{aligned} & \prod_{j=1}^n x_j^{-\lambda_j} \int_{B_1(x)} |f(y)| \left\{ \int_0^\infty \frac{e^{-\sum_{j=1}^n (x_j-y_j)^2/4t}}{t^{n+1}} dt \right\}^{1/2} \prod_{j=1}^n y_j^{\lambda_j} dy \\ & \leq C \left(\sum_{k=0}^\infty \int_{2^{k+m_1} \leq |\bar{x}-\bar{y}| < 2^{k+1+m_1}} \frac{\tilde{f}_m(\bar{y})}{(2^{2m_1-2} + |\bar{x}-\bar{y}|^2)^{\frac{n}{2}}} d\bar{y} + \int_{|\bar{x}-\bar{y}| < 2^{m_1}} \frac{\tilde{f}_m(\bar{y})}{(2^{2m_1-2} + |\bar{x}-\bar{y}|^2)^{\frac{n}{2}}} d\bar{y} \right) \\ & \leq C \sum_{k=0}^\infty \frac{1}{2^{(m_1+k)n}} \int_{|\bar{x}-\bar{y}| < 2^{k+m_1}} \tilde{f}_m(\bar{y}) d\bar{y} \\ & \leq \frac{C}{2^{m_1}} M_{n-1}(\tilde{f}_m)(\bar{x}), \end{aligned}$$

where $\bar{x} = (x_2, \dots, x_n)$ and M_{n-1} denotes the Hardy-Littlewood maximal function on $\mathbb{R}^{n-1}$.

For every $\beta > 0$, the maximal function theorem leads to

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \prod_{j=1}^n x_j^{-\lambda_j} \int_{B_1(x)} |f(y)| \left\{ \int_0^\infty \frac{e^{-\sum_{j=1}^n \frac{(x_j-y_j)^2}{4t}}}{t^{n+1}} dt \right\}^{\frac{1}{2}} \prod_{j=1}^n y_j^{\lambda_j} dy > \beta\}) \\
& \leq m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : M_{n-1}(\tilde{f}_m)(\bar{x}) > \beta 2^{m_1} M\}) \\
& \leq 2^{\sum_{j=1}^n \lambda_j m_j} m_0^{(n)}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : M_{n-1}(\tilde{f}_m)(\bar{x}) > \beta 2^{m_1} M\}) \\
& \leq 2^{\sum_{j=1}^n \lambda_j m_j} 2^{m_1} m_0^{(n-1)}(\{\bar{x} \in \mathbb{R}^{n-1} : M_{n-1}(\tilde{f}_m)(\bar{x}) > \beta 2^{m_1} M\}) \\
& \leq 2^{\sum_{j=1}^n \lambda_j m_j} \frac{C}{\beta} \|\tilde{f}_m\|_{L^1(\mathbb{R}^{n-1}, d\bar{x})} \\
& \leq 2^{\sum_{j=1}^n \lambda_j m_j} \frac{C}{\beta} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| dy \\
& \leq \frac{C}{\beta} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

Here, C and M denote positive constants that do not depend on $m \in \mathbb{Z}^n$, and $m_0^{(k)}$ represents the Lebesgue measure on $\mathbb{R}^k$.

From (15) and (17) we deduce that

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} Y_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \\
& \leq \frac{C}{\gamma} \sum_{m \in \mathbb{Z}^n} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
(18) \quad & \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)}.
\end{aligned}$$

Finally, putting together (13), (14) and (18) we conclude that

$$m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |g_{1,loc}^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma\}) \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)}.$$

Thus Claim 1 is established. $\square$

We now prove Claim 2. Minkowski inequality allow us to write

$$\begin{aligned}
G^{\lambda_1, \dots, \lambda_n}(f)(x) & \leq \sum_{i=1}^n \left\{ \int_0^\infty t \left| \int_{(0, \infty)^n} \left(\frac{\partial}{\partial t} W_t^{\lambda_i}(x_i, y_i) \prod_{j=1; j \neq i}^n W_t^{\lambda_j}(x_j, y_j) \right. \right. \right. \\
& \quad \left. \left. - \prod_{j=1}^n \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j) (x_i y_i)^{-\lambda_i} \frac{\partial}{\partial t} \left(\frac{e^{-(x_i - y_i)^2/4t}}{2\sqrt{\pi t}} \right) \right. \right. \\
& \quad \left. \left. \times \prod_{j=1; j \neq i}^n \frac{e^{-(x_j - y_j)^2/4t}}{2\sqrt{\pi t}} (x_j y_j)^{-\lambda_j} \right) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \Bigg\}^{1/2} \\
& = \sum_{i=1}^n G_i^{\lambda_1, \dots, \lambda_n}(f)(x), \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.
\end{aligned}$$

By taking into account symmetries to show Claim 2, it is sufficient to prove that $G_1^{\lambda_1, \dots, \lambda_n}$ defines a bounded operator from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Again by using Minkowski inequality we have that

$$\begin{aligned}
 G_1^{\lambda_1, \dots, \lambda_n}(f)(x) &\leq \left\{ \int_0^\infty t \left| \int_{(0, \infty)^n} \left(\frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) - \chi_{\{x_1/2 < y_1 < 2x_1\}}(y_1)(x_1 y_1)^{-\lambda_1} \right. \right. \right. \\
 &\quad \times \left. \left. \frac{\partial}{\partial t} \left(\frac{e^{-(x_1 - y_1)^2/4t}}{2\sqrt{\pi t}} \right) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \Bigg\}^{1/2} \\
 &\quad + \sum_{i=2}^n \left\{ \int_0^\infty t \left| \int_{(0, \infty)^n} \chi_{\{x_i/2 < y_i < 2x_i\}}(y_i)(x_i y_i)^{-\lambda_i} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1 - y_1)^2/4t}}{2\sqrt{\pi t}} \right) \right. \right. \\
 &\quad \times \prod_{j=2}^{i-1} \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j)(x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2/4t}}{2\sqrt{\pi t}} \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) \\
 &\quad \times \left. \left. \left(W_t^{\lambda_i}(x_i, y_i) - \chi_{\{x_i/2 < y_i < 2x_i\}}(y_i)(x_i y_i)^{-\lambda_i} \frac{e^{-(x_i - y_i)^2/4t}}{2\sqrt{\pi t}} \right) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2} \\
 (19) \quad &= \sum_{i=1}^n G_{1,i}^{\lambda_1, \dots, \lambda_n}(f)(x), \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.
 \end{aligned}$$

We are going to study $G_{1,i}^{\lambda_1, \dots, \lambda_n}$, $i = 1, \dots, n$. Firstly we consider $G_{1,1}^{\lambda_1, \dots, \lambda_n}$. By splitting the integral on $y_1 \in (0, \infty)$ we get

$$\begin{aligned}
 G_{1,1}^{\lambda_1, \dots, \lambda_n}(f)(x) &\leq \left\{ \int_0^\infty t \left| \int_0^{x_1/2} \int_{(0, \infty)^{n-1}} \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \\
 &\quad + \left\{ \int_0^\infty t \left| \int_{2x_1}^\infty \int_{(0, \infty)^{n-1}} \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \\
 &\quad + \left\{ \int_0^\infty t \left| \int_{x_1/2}^{2x_1} \int_{(0, \infty)^{n-1}} \left(\frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) - (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1 - y_1)^2/4t}}{2\sqrt{\pi t}} \right) \right) \right. \right. \\
 &\quad \times \left. \left. \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{\frac{1}{2}} \\
 (20) \quad &= \sum_{i=1}^3 G_{1,1;i}^{\lambda_1, \dots, \lambda_n}(f)(x), \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.
 \end{aligned}$$

According to [8, proof of Lemma 8] we have, for every $\lambda > -1/2$, $t, x, y \in (0, \infty)$ and $y < x/2$,

$$(21) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) \right| \leq C \frac{(xy)^{-\lambda} x^2}{t^{5/2}} e^{-x^2/16t}, \quad xy > t,$$

and

$$(22) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) \right| \leq C \frac{1}{t^{\lambda+3/2}} e^{-(x^2+y^2)/8t}, \quad xy \leq t.$$

From (21) we deduce that, when $0 < y < x/2 < \infty$,

$$(23) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) \right| \leq C \frac{(xy)^{-\lambda-1/2} x^3}{t^{5/2}} e^{-x^2/16t} \leq \frac{C}{t^{\lambda+3/2}} e^{-x^2/20t}, \quad xy > t > 0,$$

together with (22), implies that, for every $\lambda > -1/2$, $t, x, y \in (0, \infty)$, and $y < x/2$,

$$(24) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) \right| \leq \frac{C}{t^{\lambda+3/2}} e^{-x^2/20t}.$$

It will be proved that the operator defined by $G_{1,1;1}^{\lambda_1, \dots, \lambda_n}(f)(x)$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ when we show that, for every $1 \leq l \leq m \leq n$, the operator $S_{l,m}^{\lambda_1, \dots, \lambda_n}$ defined by

$$S_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) = \left\{ \int_0^\infty t \left| \int_0^{x_1/2} \dots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \dots \int_{x_m/2}^{2x_m} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty \right. \right. \\ \left. \left. \left| \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2}, \quad x \in (0, \infty)^n,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

By using (6) since the operator H_∞^k is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself, for every $k \in \mathbb{N}$ and $\alpha_j > -1/2$, $j = 1, \dots, k$, in order to prove that the operator $S_{l,m}^{\lambda_1, \dots, \lambda_n}$, $1 \leq l \leq m \leq n$, is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ it is sufficient to show that the operator

$$\mathcal{S}_{l,m}^{\alpha_1, \dots, \alpha_m}(f)(x) = \left\{ \int_0^\infty t \left| \int_0^{x_1/2} \dots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \dots \int_{x_m/2}^{2x_m} \left| \frac{\partial}{\partial t} W_t^{\alpha_1}(x_1, y_1) \right| \right. \right. \\ \left. \left. \times \prod_{j=2}^m W_t^{\alpha_j}(x_j, y_j) |f(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \right|^2 dt \right\}^{1/2}, \quad x = (x_1, \dots, x_m) \in (0, \infty)^m,$$

is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$, for every $\alpha_j > -1/2$, $j = 1, \dots, m$, and $1 \leq l \leq m$.

Let $\alpha_j > -1/2$, $j = 1, \dots, m$. According to (9) and (24), we can write

$$\mathcal{S}_{m,m}^{\alpha_1, \dots, \alpha_m}(f)(x) \leq C \left\{ \int_0^\infty t \left| \int_0^{x_1/2} \dots \int_0^{x_m/2} \frac{e^{-(\sum_{j=1}^m x_j^2)/20t}}{t^{\sum_{j=1}^m (\alpha_j+1/2)+1}} |f(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \right|^2 dt \right\}^{1/2} \\ \leq C \left\{ \int_0^\infty \frac{e^{-(\sum_{j=1}^m x_j^2)/10t}}{t^{2 \sum_{j=1}^m (\alpha_j+1/2)+1}} dt \right\}^{1/2} \int_0^{x_1/2} \dots \int_0^{x_m/2} |f(y)| \prod_{j=1}^m y_j^{2\alpha_j} dy \\ \leq CL_{\alpha_1, \dots, \alpha_m}(|f|)(x), \quad x = (x_1, \dots, x_m) \in (0, \infty)^m.$$

Then, $\mathcal{S}_{m,m}^{\alpha_1, \dots, \alpha_m}$ is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$, because $L_{\alpha_1, \dots, \alpha_m}$ has this property.

Assume that $1 \leq l < m$. By (12), since $H_{loc}^{\alpha_1, \dots, \alpha_k}$, $1 \leq k \leq m$, is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself and $\mathcal{S}_{m,m}^{\alpha_1, \dots, \alpha_m}$ from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, [14, Proposition 1] implies that $\mathcal{S}_{l,m}^{\alpha_1, \dots, \alpha_m}$ is bounded from $L^1((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^m, \prod_{j=1}^m x_j^{2\alpha_j} dx)$.

$\prod_{j=1}^m x_j^{2\alpha_j} dx$), provided that, for every $k \in \mathbb{N}$, $m < k \leq n$, the operator

$$T_{l,k}^{\alpha_1, \dots, \alpha_k}(f)(x) = \left\{ \int_0^\infty t \left| \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_k/2}^{2x_k} e^{\frac{-(\sum_{j=1}^l x_j^2 + \sum_{j=l+1}^k (x_j - y_j)^2)/20t}{\sum_{j=1}^l (\alpha_j + \frac{1}{2}) + 1 + \frac{k-l}{2}}} \right. \right. \\ \left. \left. \times \prod_{j=l+1}^k (x_j y_j)^{-\alpha_j} |f(y)| \prod_{j=1}^k y_j^{2\alpha_j} dy \right|^2 dt \right\}^{1/2}, \quad x = (x_1, \dots, x_k) \in (0, \infty)^k,$$

has this property. Let $k \in \mathbb{N}$, $m < k \leq n$. Minkowski's inequality leads to

$$T_{l,k}^{\alpha_1, \dots, \alpha_k}(f)(x) \leq C \int_0^{x_1/2} \cdots \int_0^{x_l/2} \int_{x_{l+1}/2}^{2x_{l+1}} \cdots \int_{x_k/2}^{2x_k} \prod_{j=l+1}^k (x_j y_j)^{-\alpha_j} |f(y)| \\ \times \left\{ \int_0^\infty \frac{e^{\frac{-(\sum_{j=1}^l x_j^2 + \sum_{j=l+1}^k (x_j - y_j)^2)/10t}{2 \sum_{j=1}^l (\alpha_j + \frac{1}{2}) + 1 + k - l}} dt \right\}^{1/2} \prod_{j=1}^k y_j^{2\alpha_j} dy \\ \leq C \mathcal{H}_{l,k}^{\alpha_1, \dots, \alpha_k}(|f|)(x), \quad x = (x_1, \dots, x_k) \in (0, \infty)^k.$$

Hence, $T_{l,k}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, because $\mathcal{H}_{l,k}^{\alpha_1, \dots, \alpha_k}$ has this property.

We have proved that $G_{1,1;1}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

In order to see that $G_{1,1;2}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, we must prove that, for every $1 < l \leq m \leq n$, the operator

$$S_{l,m}^{\alpha_1, \dots, \alpha_m}(f)(x) = \left\{ \int_0^\infty t \left| \int_{2x_1}^\infty \cdots \int_{2x_l}^\infty \int_0^{x_{l+1}/2} \cdots \int_0^{x_m/2} \int_{x_{m+1}/2}^{2x_{m+1}} \cdots \int_{x_n/2}^{2x_n} \right. \right. \\ \left. \left. \left| \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2}, \quad x \in (0, \infty)^m,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Note firstly that if $l < m$, we can proceed as in the previous cases (in the analysis of $G_{1,1;1}^{\lambda_1, \dots, \lambda_n}$) to see that $S_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Suppose now that $l = m$. Since the operator H_∞^k is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself, for every $\alpha_j > -1/2$, $j = 1, \dots, k$, in order to prove that $S_{m,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, it is sufficient to show that the operator

$$S_k^{\alpha_1, \dots, \alpha_k}(f)(x) = \left\{ \int_0^\infty t \left| \int_{2x_1}^\infty \int_{\frac{x_2}{2}}^{2x_2} \cdots \int_{\frac{x_k}{2}}^{2x_k} \left| \frac{\partial}{\partial t} W_t^{\alpha_1}(x_1, y_1) \right| \prod_{j=2}^k W_t^{\alpha_j}(x_j, y_j) |f(y)| \prod_{j=1}^k y_j^{2\alpha_j} dy \right|^2 dt \right\}^{\frac{1}{2}}$$

is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, for every $\alpha_j > -1/2$, $j = 1, \dots, k$, and for each $x = (x_1, \dots, x_k) \in (0, \infty)^k$.

If $\alpha_j > -1/2$, $j = 1, \dots, k$, since $H_{loc}^{\beta_1, \dots, \beta_l}$ is bounded from $L^1((0, \infty)^l, \prod_{j=1}^l x_j^{2\beta_j} dx)$ into itself when $\beta_j > -1/2$, $j = 1, \dots, l$, and $l \in \mathbb{N}$, by taking into account that $W_t^\lambda(x, y) = W_t^\lambda(y, x)$, $t, x, y \in (0, \infty)$, and (12) and (24), the desired property for $S_k^{\alpha_1, \dots, \alpha_k}$ follows when we see that the operator

$$T_k^{\beta_1, \dots, \beta_l}(f)(x) = \left\{ \int_0^\infty t \left| \int_{2x_1}^\infty \int_{x_2/2}^{2x_2} \dots \int_{x_l/2}^{2x_l} \frac{e^{-\frac{y_1^2 + \sum_{j=2}^l (x_j - y_j)^2}{20t}}}{t^{\beta_1 + \frac{3}{2} + \frac{l-1}{2}}} \prod_{j=2}^l (x_j y_j)^{-\beta_j} f(y) \prod_{j=1}^l y_j^{2\beta_j} dy \right|^2 dt \right\}^{\frac{1}{2}}$$

is bounded from $L^1((0, \infty)^l, \prod_{j=1}^l x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^l, \prod_{j=1}^l x_j^{2\beta_j} dx)$, for every $\beta_j > -1/2$, $j = 1, \dots, l$, $l \in \mathbb{N}$, and $x = (x_1, \dots, x_l) \in (0, \infty)^l$.

Let $\beta_j > -1/2$, $j = 1, \dots, l$ and $l \in \mathbb{N}$. Minkowski's inequality leads to

$$\begin{aligned} |T_k^{\beta_1, \dots, \beta_l}(f)(x)| &\leq \int_{2x_1}^\infty \int_{x_2/2}^{2x_2} \dots \int_{x_l/2}^{2x_l} \left\{ \int_0^\infty \frac{e^{-\frac{y_1^2 + \sum_{j=2}^l (x_j - y_j)^2}{10t}}}{t^{2\beta_1 + l + 1}} dt \right\}^{\frac{1}{2}} \prod_{j=2}^l (x_j y_j)^{-\beta_j} |f(y)| \prod_{j=1}^l y_j^{2\beta_j} dy \\ &\leq C \int_{2x_1}^\infty \int_{x_2/2}^{2x_2} \dots \int_{x_l/2}^{2x_l} \frac{\prod_{j=2}^l (x_j y_j)^{-\beta_j} |f(y)|}{(y_1^2 + \sum_{j=2}^l (x_j - y_j)^2)^{\beta_1 + \frac{l}{2}}} \prod_{j=1}^l y_j^{2\beta_j} dy, \quad x \in (0, \infty)^l. \end{aligned}$$

Then, by proceeding as in the proof of the main property (Case 3) in [25] we can see that the operator $T_k^{\beta_1, \dots, \beta_l}$ is bounded from $L^1((0, \infty)^l, \prod_{j=1}^l x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^l, \prod_{j=1}^l x_j^{2\beta_j} dx)$.

We conclude that $G_{1,1;2}^{\lambda_1, \dots, \lambda_n}$ defines a bounded operator from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

We are going to work with $G_{1,1;3}^{\lambda_1, \dots, \lambda_n}$. According to [8, proof of Lemma 8] we have, for every $\lambda > -1/2$, that

$$(25) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) - (xy)^{-\lambda} \frac{\partial}{\partial t} \mathbb{W}_t(x, y) \right| \leq C \frac{(xy)^{-\lambda-1}}{\sqrt{t}} e^{-(x-y)^2/8t}, \quad t, x, y \in (0, \infty), \quad xy \geq t,$$

and

$$(26) \quad \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) \right| \leq C \frac{e^{-(x-y)^2/8t}}{t^{\lambda+3/2}}, \quad t, x, y \in (0, \infty), \quad xy < t.$$

By combining (16), (25) and (26) in [[8], Lemma 8] it was established that

$$(27) \quad \left(\int_0^\infty t \left| \frac{\partial}{\partial t} W_t^\lambda(x, y) - (xy)^{-\lambda} \frac{\partial}{\partial t} \mathbb{W}_t(x, y) \right|^2 dt \right)^{\frac{1}{2}} \leq \frac{C}{x^{2\lambda+1}}, \quad 0 < \frac{x}{2} < y < 2x,$$

provided that $\lambda > -1/2$.

The operator defined by $G_{1,1;3}^{\lambda_1,\dots,\lambda_n}$ is bounded from $L^1((0,\infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0,\infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ provided that, for every $1 \leq l \leq m \leq n$ and $\alpha_j > -1/2$, $j = 1, \dots, n$, the operator

$$S_{l,m}^{\alpha_1,\dots,\alpha_n}(f)(x) = \left\{ \int_0^\infty t \left| \int_{x_1/2}^{2x_1} \left| \frac{\partial}{\partial t} W_t^{\alpha_1}(x_1, y_1) - (x_1 y_1)^{-\alpha_1} \frac{\partial}{\partial t} \mathbb{W}_t(x_1, y_1) \right| \int_{\frac{x_2}{2}}^{2x_2} \dots \int_{\frac{x_l}{2}}^{2x_l} \right. \right. \\ \left. \left. \int_0^{\frac{x_{l+1}}{2}} \dots \int_0^{\frac{x_m}{2}} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty \prod_{j=2}^n W_t^{\alpha_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\alpha_j} dy \right|^2 dt \right\}^{1/2}, \quad x \in (0, \infty)^n,$$

is bounded from $L^1((0,\infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0,\infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$.

Suppose that $l < m$ and $\alpha_j > -1/2$, $j = 1, \dots, n$. According to (27) and by using Minkowski inequality we obtain that

$$S_{l,m}^{\alpha_1,\dots,\alpha_n}(f)(x) \leq C \int_{\frac{x_1}{2}}^{2x_1} \frac{1}{x_1^{2\alpha_1+1}} \int_{\frac{x_2}{2}}^{2x_2} \dots \int_{\frac{x_l}{2}}^{2x_l} \int_0^{\frac{x_{l+1}}{2}} \dots \int_0^{\frac{x_m}{2}} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty \\ \sup_{t>0} \prod_{j=2}^n W_t^{\alpha_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\alpha_j} dy, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.$$

In Section 2 it was proved that the operator

$$T_{l,m}^{\alpha_2,\dots,\alpha_n}(g)(x) = \int_{\frac{x_2}{2}}^{2x_2} \dots \int_{\frac{x_l}{2}}^{2x_l} \int_0^{\frac{x_{l+1}}{2}} \dots \int_0^{\frac{x_m}{2}} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty \\ \sup_{t>0} \prod_{j=2}^n W_t^{\alpha_j}(x_j, y_j) |g(y)| \prod_{j=2}^n y_j^{2\alpha_j} dy, \quad x = (x_2, \dots, x_n) \in (0, \infty)^{n-1},$$

is bounded from $L^1((0,\infty)^{n-1}, \prod_{j=2}^n x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0,\infty)^{n-1}, \prod_{j=2}^n x_j^{2\alpha_j} dx)$. Moreover, $H_{loc}^{\alpha_1}$ is bounded from $L^1((0,\infty), x^{2\alpha_1} dx)$ into itself. Then, [14, Proposition 1] implies that $S_{l,m}^{\alpha_1,\dots,\alpha_n}$ is a bounded operator from $L^1((0,\infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0,\infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$.

Since the operator H_∞^k is bounded from $L^1((0,\infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $\beta_j > -1/2$, $j = 1, \dots, k$, we need only to see that the operator

$$\mathcal{S}_k^{\beta_1,\dots,\beta_k}(g)(x) = \left\{ \int_0^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \int_{\frac{x_2}{2}}^{2x_2} \dots \int_{\frac{x_k}{2}}^{2x_k} \left(\frac{\partial}{\partial t} W_t^{\beta_1}(x_1, y_1) - (x_1 y_1)^{-\beta_1} \frac{\partial}{\partial t} \mathbb{W}_t(x_1, y_1) \right) \right. \right. \\ \left. \left. \times \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|^2 dt \right\}^{1/2}, \quad x = (x_1, \dots, x_k) \in (0, \infty)^k,$$

is bounded from $L^1((0,\infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0,\infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$, when $\beta_j > -1/2$, $j = 1, \dots, k$, to show that the operator $S_{m,m}^{\alpha_1,\dots,\alpha_n}$, $m = 1, \dots, n$, has desired boundedness property.

Let $\beta_j > -1/2$, $j = 1, \dots, k$. By (16), (25) and (26) we have that

$$\mathcal{S}_k^{\beta_1,\dots,\beta_k}(g)(x) = \left\{ \left(\int_0^{x_1^2} + \int_{x_1^2}^\infty \right) t \left| \int_{\frac{x_1}{2}}^{2x_1} \int_{\frac{x_2}{2}}^{2x_2} \dots \int_{\frac{x_k}{2}}^{2x_k} \left(\frac{\partial}{\partial t} W_t^{\beta_1}(x_1, y_1) \right. \right. \right. \\ \left. \left. \left. - (x_1 y_1)^{-\beta_1} \frac{\partial}{\partial t} \mathbb{W}_t(x_1, y_1) \right) \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|^2 dt \right\}^{1/2}$$

$$\begin{aligned}
&\leq C \left[\left\{ \int_0^{x_1^2} t \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} (x_1 y_1)^{-\beta_1-1} \frac{e^{-(x_1-y_1)^2/8t}}{\sqrt{t}} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|^2 dt \right\}^{1/2} \right. \\
&\quad + \left\{ \int_{x_1^2}^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} \frac{e^{-(x_1-y_1)^2/8t}}{t^{\beta_1+3/2}} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|^2 dt \right\}^{1/2} \\
&\quad + \left. \left\{ \int_{x_1^2}^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} (x_1 y_1)^{-\beta_1} \frac{\partial}{\partial t} \mathbb{W}_t(x_1, y_1) \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|^2 dt \right\}^{1/2} \right] \\
&\leq C \left[\left\{ \int_0^{x_1^2} t dt \right\}^{1/2} \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} (x_1 y_1)^{-\beta_1-1} \frac{e^{-(x_1-y_1)^2/8t}}{\sqrt{t}} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy \right. \\
&\quad + \left\{ \int_{x_1^2}^\infty \frac{1}{t^{2\beta_1+2}} dt \right\}^{1/2} \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy \\
&\quad + \left. \left\{ \int_{x_1^2}^\infty \frac{dt}{t^2} \right\}^{1/2} x_1^{-2\beta_1} \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy \right] \\
&\leq C \left[\sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} (x_1 y_1)^{-\beta_1} \frac{e^{-(x_1-y_1)^2/8t}}{\sqrt{t}} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy \right. \\
&\quad + \left. \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} \prod_{j=2}^k W_t^{\beta_j}(x_j, y_j) \frac{1}{x_1^{2\beta_1+1}} \int_{\frac{x_1}{2}}^{2x_1} g(y) y_1^{2\beta_1} dy_1 \prod_{j=2}^k y_j^{2\beta_j} dy_k \dots dy_2 \right], \quad x \in (0, \infty)^k.
\end{aligned}$$

Since the operator $H_{loc}^{\beta_1}$ is bounded from $L^1((0, \infty), y^{2\beta_1} dy)$ into itself, by taking into account the results established in the proof of Theorem 1.1 (see Section 2), we conclude that the operator $\mathcal{S}_k^{\beta_1, \dots, \beta_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$.

Thus, we prove that $G_{1,1;3}^{\lambda_1, \dots, \lambda_n}$ defines a bounded operator from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

According to (20) we establish that the operator defined by $G_{1,1}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Assume now that $i = 2, \dots, n$. We analyze the operator $G_{1,i}^{\lambda_1, \dots, \lambda_n}$. In order to do this we write

$$\begin{aligned}
G_{1,i}^{\lambda_1, \dots, \lambda_n}(f)(x) &\leq \left\{ \int_0^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_{i-1}}{2}}^{2x_{i-1}} (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{\frac{-(x_1-y_1)^2}{4t}}}{2\sqrt{\pi t}} \right) \prod_{j=2}^{i-1} (x_j y_j)^{-\lambda_j} \frac{e^{\frac{-(x_j-y_j)^2}{4t}}}{2\sqrt{\pi t}} \right. \right. \\
&\quad \times \left. \left. \int_0^{\frac{x_i}{2}} \int_{(0, \infty)^{n-i}} \prod_{j=i}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2} + \left\{ \int_0^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_{i-1}}{2}}^{2x_{i-1}} (x_1 y_1)^{-\lambda_1} \right. \right. \\
&\quad \times \left. \left. \frac{\partial}{\partial t} \left(\frac{e^{\frac{-(x_1-y_1)^2}{4t}}}{2\sqrt{\pi t}} \right) \prod_{j=2}^{i-1} (x_j y_j)^{-\lambda_j} \frac{e^{\frac{-(x_j-y_j)^2}{4t}}}{2\sqrt{\pi t}} \int_{2x_i}^\infty \int_{(0, \infty)^{n-i}} \prod_{j=i}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2}
\end{aligned}$$

$$\begin{aligned}
& + \left\{ \int_0^\infty t \left| \int_{\frac{x_1}{2}}^{2x_1} \cdots \int_{\frac{x_{i-1}}{2}}^{2x_{i-1}} (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1 - y_1)^2}{4t}}}{2\sqrt{\pi t}} \right) \prod_{j=2}^{i-1} (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{2\sqrt{\pi t}} \right. \right. \\
& \times \left. \left. \int_{\frac{x_i}{2}}^{2x_i} \left(W_t^{\lambda_i}(x_i, y_i) - (x_i y_i)^{-\lambda_i} \frac{e^{-\frac{(x_i - y_i)^2}{4t}}}{2\sqrt{\pi t}} \right) \int_{(0, \infty)^{n-i}} \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|^2 dt \right\}^{1/2} \\
& = \sum_{l=1}^3 G_{1,i;l}(f)(x), \quad x = (x_1, \dots, x_n) \in (0, \infty)^n.
\end{aligned}$$

In order to analyze the boundedness property for $G_{1,i;1}$ and $G_{1,i;2}$ we can follow a similar way to the one employed to study $G_{1,1;1}$ and $G_{1,1;2}$, respectively.

On the other hand, by (3) we get, for every $\lambda > -1/2$, $t, x, y \in (0, \infty)$,

$$(28) \quad |W_t^\lambda(x, y) - (xy)^{-\lambda} \mathbb{W}_t(x, y)| \leq C \left(\frac{1}{t^{\lambda+1/2}} + \frac{(xy)^{-\lambda}}{\sqrt{t}} \right) e^{-\frac{x^2+y^2}{4t}}, \quad xy \leq t.$$

Also, (4) implies that

$$(29) \quad |W_t^\lambda(x, y) - (xy)^{-\lambda} \mathbb{W}_t(x, y)| \leq C(xy)^{-\lambda-1} \sqrt{t} e^{-\frac{(x-y)^2}{4t}}, \quad xy > t,$$

when $\lambda > -1/2$, $t, x, y \in (0, \infty)$.

By using (28) and (29), and proceeding as in the study of $G_{1,1;3}$ we can prove that $G_{1,i;3}$ defines a bounded operator from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Thus, we conclude that $G_{1,i}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Hence, the operator G_1 is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, and the proof of Claim 2 is finished. $\square$

4. PROOF OF THEOREM 1.3

In this section we prove that the Riesz transforms $R_i^{\lambda_1, \dots, \lambda_n}$, $i = 1, \dots, n$, associated with the Bessel operator $\Delta_{\lambda_1, \dots, \lambda_n}$, are bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

For every $i = 1, \dots, n$, and $\lambda_j > -1/2$, $j = 1, \dots, n$, the Riesz transform $R_i^{\lambda_1, \dots, \lambda_n}$ is formally defined by

$$(30) \quad R_i^{\lambda_1, \dots, \lambda_n} = \frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2},$$

where $\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2}$ denotes the negative square root of the operator $\Delta_{\lambda_1, \dots, \lambda_n}$. We are going to precise the definition (30).

Assume that $0 < \beta < \sum_{j=1}^n (\lambda_j + 1/2) + 1$ and $\lambda_j > -1/2$, $j = 1, \dots, n$. We define the negative power $\Delta_{\lambda_1, \dots, \lambda_n}^{-\beta}$ on $C_c^\infty((0, \infty)^n)$ as follows

$$\Delta_{\lambda_1, \dots, \lambda_n}^{-\beta} f(x) = \frac{1}{\Gamma(\beta)} \int_0^\infty \left(W_t^{\lambda_1, \dots, \lambda_n}(f)(x) - \chi_{(1, \infty)}(t) \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right)$$

$$\times \int_{(0,\infty)^n} f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \Big) t^{\beta-1} dt, \quad x \in (0,\infty)^n,$$

where $f \in C_c^\infty((0,\infty)^n)$.

Let $f \in C_c^\infty((0,\infty)^n)$. We can write

$$(31) \quad \Delta_{\lambda_1, \dots, \lambda_n}^{-\beta} f(x) = \int_{(0,\infty)^n} f(y) K_{\beta}^{\lambda_1, \dots, \lambda_n}(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0,\infty)^n,$$

being, for each $x, y \in (0,\infty)^n$, $x \neq y$,

$$K_{\beta}^{\lambda_1, \dots, \lambda_n}(x, y) = \frac{1}{\Gamma(\beta)} \int_0^\infty \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \chi_{(1,\infty)}(t) \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) t^{\beta-1} dt.$$

In order to prove the integral representation (31) for $\Delta_{\lambda_1, \dots, \lambda_n}^{-\beta}$ it is sufficient to show that, for every $x \in (0,\infty)^n$,

$$\int_{(0,\infty)^n} |f(y)| \int_0^\infty \left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \chi_{(1,\infty)}(t) \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right| t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy < \infty.$$

Indeed, let $x = (x_1, \dots, x_n) \in (0,\infty)^n$. According to (3) and (4) and denoting $K = \text{supp} f$, we have that

$$\begin{aligned} & \int_{(0,\infty)^n} |f(y)| \int_0^1 \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\ & \leq C \int_K \int_0^1 t^{\beta-1-n/2} e^{-\sum_{j=1}^n (x_j - y_j)^2 / 4t} dt dy \\ & \leq C \int_K \frac{1}{|x - y|^{n-2\beta}} dy < \infty, \end{aligned}$$

because $\beta > 0$. In the last inequality we have used [32, Lemma 1.1].

On the other hand, by using again (3), (4) and (5) it follows

$$\begin{aligned} & \left| W_t^\lambda(x, y) - \frac{t^{-\lambda-1/2}}{2^{2\lambda} \Gamma(\lambda + 1/2)} \right| = \left| \frac{1}{(2t)^{\lambda+1/2}} \left(\frac{xy}{2t} \right)^{-\lambda+1/2} I_{\lambda-1/2} \left(\frac{xy}{2t} \right) e^{-\frac{x^2+y^2}{4t}} - \frac{t^{-\lambda-1/2}}{2^{2\lambda} \Gamma(\lambda + 1/2)} \right| \\ & \leq \frac{1}{(2t)^{\lambda+1/2}} e^{-\frac{x^2+y^2}{4t}} \left| \left(\frac{xy}{2t} \right)^{-\lambda+1/2} I_{\lambda-1/2} \left(\frac{xy}{2t} \right) - \frac{1}{2^{\lambda-1/2} \Gamma(\lambda + 1/2)} \right| \\ & \quad + \frac{t^{-\lambda-1/2}}{2^{2\lambda} \Gamma(\lambda + 1/2)} \left| e^{-\frac{x^2+y^2}{4t}} - 1 \right| \\ & \leq \frac{xy}{(2t)^{\lambda+3/2}} e^{-\frac{x^2+y^2}{4t}} \sup_{z \in (0, \frac{xy}{2t})} \left| \frac{d}{dz} \left(z^{-\lambda+1/2} I_{\lambda-1/2}(z) \right) \right| + C \frac{x^2 + y^2}{t^{\lambda+3/2}} \\ & \leq C \left(\frac{(xy)^2}{t^{\lambda+5/2}} e^{-\frac{(x-y)^2}{4t}} + \frac{x^2 + y^2}{t^{\lambda+3/2}} \right) \\ (32) \quad & \leq C \frac{x^2 + y^2}{t^{\lambda+3/2}} \left(\frac{xy}{t} + 1 \right), \quad t, x, y \in (0, \infty). \end{aligned}$$

Then, by (3) we get

$$\begin{aligned}
& \int_{(0,\infty)^n} |f(y)| \int_1^\infty \left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right| t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& \leq C \sum_{i=1}^n \int_K \int_1^\infty \prod_{j=1}^{i-1} \frac{t^{-(\lambda_j+1/2)}}{2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \left| W_t^{\lambda_i}(x_i, y_i) - \frac{t^{-(\lambda_i+1/2)}}{2^{2\lambda_i} \Gamma(\lambda_i + 1/2)} \right| \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) t^{\beta-1} dt dy \\
& \leq C \int_K \int_1^\infty t^{-\sum_{j=1}^n (\lambda_j+1/2)-2+\beta} dt dy < \infty,
\end{aligned}$$

because $\beta < \sum_{j=1}^n (\lambda_j + 1/2) + 1$.

In particular, we have

$$\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) = \int_{(0,\infty)^n} f(y) K_{1/2}^{\lambda_1, \dots, \lambda_n}(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty).$$

Remark 4.1. Note that if $0 < \beta < \sum_{j=1}^n (\lambda_j + 1/2)$, for every $f \in C_c^\infty((0, \infty)^n)$,

$$\int_{(0,\infty)^n} |f(y)| \int_0^\infty \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy < \infty, \quad x \in (0, \infty)^n,$$

and we can define the function

$$(\Delta_{\lambda_1, \dots, \lambda_n})^{-\beta} f(x) = \int_{(0,\infty)^n} \frac{f(y)}{\Gamma(\beta)} \int_0^\infty \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n,$$

when $f \in C_c^\infty((0, \infty)^n)$. Then, if $0 < \beta < \sum_{j=1}^n (\lambda_j + 1/2)$ and $f \in C_c^\infty((0, \infty)^n)$, for each $x \in (0, \infty)^n$,

$$\Delta_{\lambda_1, \dots, \lambda_n}^{-\beta} f(x) - (\Delta_{\lambda_1, \dots, \lambda_n})^{-\beta} f(x) = \frac{-1}{\Gamma(\beta)} \int_{(0,\infty)^n} f(y) \int_1^\infty \prod_{j=1}^n \frac{t^{-(\lambda_j+1/2)}}{2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} t^{\beta-1} dt \prod_{j=1}^n y_j^{2\lambda_j} dy$$

and, for every $i = 1, \dots, n$,

$$\frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-\beta} f(x) = \frac{\partial}{\partial x_i} (\Delta_{\lambda_1, \dots, \lambda_n})^{-\beta} f(x),$$

provided that this derivative exists.

For every $0 < \beta < n/2$, the fractional power $\Delta^{-\beta}$ is defined on $C_c^\infty(\mathbb{R}^n)$ by

$$\begin{aligned}
\Delta^{-\beta} f(x) &= \frac{1}{\Gamma(\beta)} \int_0^\infty t^{\beta-1} \frac{1}{(2\sqrt{\pi})^n} \int_{\mathbb{R}^n} \frac{e^{-|x-y|^2/4t}}{t^{n/2}} f(y) dy dt \\
&= \int_{\mathbb{R}^n} f(y) \frac{1}{\Gamma(\beta)(2\sqrt{\pi})^n} \int_0^\infty t^{\beta-1-n/2} e^{-|x-y|^2/4t} dt dy \\
&= \frac{\Gamma(n/2 - \beta)}{\pi^{n/2} 4^\beta \Gamma(\beta)} \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-2\beta}} dy, \quad x \in \mathbb{R}^n \text{ and } f \in C_c^\infty(\mathbb{R}^n).
\end{aligned}$$

Note that all the above integrals are absolutely convergent. In particular, since $n > 1$, we have that, for every $f \in C_c^\infty(\mathbb{R}^n)$,

$$\Delta^{-1/2} f(x) = \frac{1}{2^n \pi^{\frac{n+1}{2}}} \int_{\mathbb{R}^n} f(y) \int_0^\infty \frac{e^{-|x-y|^2/4t}}{t^{\frac{n+1}{2}}} dt dy, \quad x \in \mathbb{R}^n.$$

A crucial result to prove Theorem 1.3 is the following one.

Proposition 4.1. *Let $f \in C_c^\infty((0, \infty)^n)$. Assume that $\lambda_j > -1/2$, $j = 1, \dots, n$. Then, for every $i = 1, \dots, n$, it has*

$$(33) \quad \frac{\partial}{\partial x_i} \left(\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) - \prod_{j=1}^n x_j^{-\lambda_j} \Delta^{-1/2} \left(\prod_{j=1}^n y_j^{\lambda_j} f \right)(x) \right) \\ = \int_{(0, \infty)^n} f(y) \left(R_i^{\lambda_1, \dots, \lambda_n}(x, y) - \mathcal{R}_i^{\lambda_1, \dots, \lambda_n}(x, y) \right) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad a.e. \ x \in (0, \infty)^n,$$

where

$$\mathcal{R}_i^{\lambda_1, \dots, \lambda_n}(x, y) = \frac{1}{2^n \pi^{\frac{n+1}{2}}} \frac{\partial}{\partial x_i} \left(\int_0^\infty \frac{e^{-|x-y|^2/4t}}{t^{\frac{n+1}{2}}} dt \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \right), \quad x, y \in \mathbb{R}^n, \ x \neq y,$$

and

$$R_i^{\lambda_1, \dots, \lambda_n}(x, y) = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{\partial}{\partial x_i} W_t^{\lambda_i}(x_i, y_i) \prod_{j=1, j \neq i}^n W_t^{\lambda_j}(x_j, y_j) \frac{dt}{\sqrt{t}}, \quad x, y \in (0, \infty)^n, \ x \neq y.$$

Moreover, the integral in (33) is absolutely convergent.

Proof. We are going to prove (33) for $i = 1$. We denote $K = \text{supp} f$. Suppose that Ω is a compact subset of $(0, \infty)^n$. There exist $0 < a < 1$ and $b > 1$, such that for every $y \in K$ and $x \in \Omega$, $\frac{x_j y_j}{t} \geq 1$, when $0 < t < a$, and $\frac{x_j y_j}{t} \leq 1$, when $b < t < \infty$, for every $j = 1, \dots, n$. We can write, for each $x, y \in (0, \infty)^n$,

$$(34) \quad K_{1/2}^{\lambda_1, \dots, \lambda_n}(x, y) - \frac{\prod_{j=1}^n (x_j y_j)^{-\lambda_j}}{2^n \pi^{\frac{n+1}{2}}} \int_0^\infty \frac{e^{-|x-y|^2/4t}}{t^{\frac{n+1}{2}}} dt \\ = \frac{1}{\sqrt{\pi}} \left\{ \int_0^a \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2/4t}}{2\sqrt{\pi t}} \right) \frac{dt}{\sqrt{t}} \right. \\ + \int_a^1 \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2/4t}}{2\sqrt{\pi t}} \right) \frac{dt}{\sqrt{t}} \\ + \int_1^b \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \frac{\prod_{j=1}^n (x_j y_j)^{-\lambda_j}}{2^n \pi^{\frac{n}{2}}} \frac{e^{-|x-y|^2/4t}}{t^{\frac{n}{2}}} \right) \frac{dt}{\sqrt{t}} \\ \left. + \int_b^\infty \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \frac{\prod_{j=1}^n (x_j y_j)^{-\lambda_j}}{2^n \pi^{\frac{n}{2}}} \frac{e^{-|x-y|^2/4t}}{t^{\frac{n}{2}}} \right) \frac{dt}{\sqrt{t}} \right\}.$$

According to (4), we have for every $x, y \in (0, \infty)^n$ and $t > 0$,

$$\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) = \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{\sqrt{2t}} \left(\frac{x_j y_j}{2t} \right)^{1/2} I_{\lambda_j - 1/2} \left(\frac{x_j y_j}{2t} \right) e^{-(x_j^2 + y_j^2)/4t} \\ = \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \left(\sum_{k=0}^n (-1)^k [\lambda_j - 1/2, k] \left(\frac{t}{x_j y_j} \right)^k + f_n^{\lambda_j - 1/2} \left(\frac{x_j y_j}{2t} \right) \right),$$

where, for every $\nu > -1$, f_n^ν is a $C^\infty(0, \infty)$ -function such that $f_n^\nu(z) = O(z^{-(n+1)})$, as $z \rightarrow \infty$. Then, for each $x, y \in (0, \infty)^n$ and $t > 0$,

$$(35) \quad \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) = \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \times \left(1 + \sum_{\substack{k_1, \dots, k_n \in \mathbb{N} \\ k_1, \dots, k_n \leq n \\ (k_1, \dots, k_n) \neq 0}} c_{k_1, \dots, k_n} \frac{t^{k_1 + \dots + k_n}}{\prod_{j=1}^n (x_j y_j)^{k_j}} + g_n(x, y, t) \right),$$

where $c_{k_1, \dots, k_n} \in \mathbb{R}$, $k_1, \dots, k_n \in \mathbb{N}$, $k_1, \dots, k_n \leq n$, and $(k_1, \dots, k_n) \neq 0$, and $g_n \in C^\infty((0, \infty)^n \times (0, \infty)^n \times (0, \infty))$ and

$$|g_n(x, y, t)| \leq Ct^{n+1}, \quad t \in (0, a), \quad x \in \Omega \text{ and } y \in K.$$

Let $\nu > -1$. By (4) and (5) we get, for each $z \in (0, \infty)$,

$$(36) \quad \begin{aligned} \frac{d}{dz}(e^{-z} z^{-\nu} I_\nu(z)) &= e^{-z} z^{-\nu-1/2} (\sqrt{z} I_{\nu+1} - \sqrt{z} I_\nu(z)) \\ &= \frac{z^{-\nu-1/2}}{\sqrt{2\pi}} \left(\sum_{k=0}^{n+1} (-1)^k ([\nu+1, k] - [\nu, k]) (2z)^{-k} + f_{n+1}^{\nu+1}(z) - f_{n+1}^\nu(z) \right) \\ &= -\frac{z^{-\nu-1/2}}{\sqrt{2\pi}} \left(\sum_{k=0}^n (-1)^k ([\nu+1, k+1] - [\nu, k+1]) (2z)^{-k-1} + f_{n+1}^{\nu+1}(z) - f_{n+1}^\nu(z) \right), \end{aligned}$$

and

$$(37) \quad \begin{aligned} \frac{d}{dz}(e^{-z} z^{-\nu} I_\nu(z)) &= \frac{d}{dz} \left(\frac{z^{-\nu-1/2}}{\sqrt{2\pi}} \left(\sum_{k=0}^n (-1)^k [\nu, k] (2z)^{-k} + f_n^\nu(z) \right) \right) \\ &= -\frac{z^{-\nu-1/2}}{\sqrt{2\pi}} \left(\sum_{k=0}^n (-1)^k [\nu, k] (2\nu + 2k + 1) (2z)^{-k-1} + \frac{\nu+1/2}{z} f_n^\nu(z) - \frac{d}{dz} f_n^\nu(z) \right). \end{aligned}$$

Moreover, $[\nu+1, k+1] - [\nu, k+1] = [\nu, k](2\nu + 2k + 1)$, for $k \in \mathbb{N}$, $k \geq 1$. Indeed, let $k \geq 1$. We have that

$$\begin{aligned} [\nu+1, k+1] - [\nu, k+1] &= \frac{1}{2^{2k+2}\Gamma(k+2)} \left\{ \prod_{j=0}^k (4(\nu+1)^2 - (2j+1)^2) - \prod_{j=0}^k (4\nu^2 - (2j+1)^2) \right\} \\ &= \frac{1}{2^{2k+2}\Gamma(k+2)} \left\{ \prod_{j=0}^k (2\nu + 2(j+1) + 1)(2\nu - 2(j-1) - 1) - \prod_{j=0}^k (2\nu + 2j + 1)(2\nu - 2j - 1) \right\} \\ &= \frac{(2\nu + 2k + 3) - (2\nu - 2k - 1)}{4(k+1)2^{2k}\Gamma(k+1)} \left\{ \prod_{j=0}^{k-1} (2\nu + 2j + 1)(2\nu - 2j - 1) \right\} (2\nu + 2k + 1) \\ &= [\nu, k](2\nu + 2k + 1). \end{aligned}$$

Then, from (36) and (37) we deduce that

$$\frac{d}{dz} f_n^\nu = \frac{\nu+1/2}{z} f_n^\nu - f_{n+1}^{\nu+1} + f_{n+1}^\nu.$$

Hence,

$$(38) \quad \frac{d}{dz} f_n^\nu(z) = O\left(\frac{1}{z^{n+2}}\right), \quad \text{as } z \rightarrow \infty,$$

and we conclude that

$$(39) \quad \left| \frac{\partial}{\partial x_1} g_n(x, y, t) \right| \leq Ct^{n+1}, \quad t \in (0, a), \quad x \in \Omega \text{ and } y \in K.$$

If $k \in \mathbb{N}$, according to [32, Lemma 1.1] we get

$$(40) \quad \int_0^a e^{|x-y|^2/4t} t^{k-\frac{n+1}{2}} dt \leq \int_0^a e^{|x-y|^2/4t} t^{-\frac{n-1}{2}} dt \leq C \frac{1}{|x-y|^{n-2}}, \quad x \neq y, \quad x, y \in (0, \infty)^n.$$

Suppose that $k_1, \dots, k_n \in \mathbb{N}$ such that $k_1 + \dots + k_n \leq n$ and $k_i \geq 1$, for some $i = 1, \dots, n$. From (40) it deduces that the functions

$$h_{k_1, \dots, k_n}(z) = \int_0^a \frac{e^{-|z|^2/4t}}{t^{\frac{n+1}{2} - \sum_{j=1}^n k_j}} dt, \quad H_{k_1, \dots, k_n}(z) = \int_0^a \frac{e^{-|z|^2/4t}}{t^{\frac{n}{2}+1 - \sum_{j=1}^n k_j}} dt, \quad z \in \mathbb{R}^n \setminus \{0\}$$

are in $L^1(\Lambda)$ for every compact subset $\Lambda \subset \mathbb{R}^n$. Moreover,

$$\frac{\partial}{\partial x_1} h_{k_1, \dots, k_n}(x) = \int_0^a \frac{\partial}{\partial x_1} e^{-|x|^2/4t} t^{-\frac{n+1}{2} + \sum_{j=1}^n k_j} dt, \quad x \in \mathbb{R}^n \setminus \{0\}.$$

Since $f \in C_c^\infty((0, \infty)^n)$ by defining $f(y) = 0$, $y \in \mathbb{R}^n \setminus (0, \infty)^n$, the function

$$G_{k_1, \dots, k_n}(x) = \int_{\mathbb{R}^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n,$$

is derivable with respect to x_1 on $(0, \infty)^n$, and

$$\begin{aligned} \frac{\partial}{\partial x_1} G_{k_1, \dots, k_n}(x) &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad + \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j (x_j - y_j))^{-\lambda_j - k_j} \frac{\partial}{\partial x_1} [(x_1 - y_1)^{\lambda_1 - k_1} f(x - y)] h_{k_1, \dots, k_n}(y) \prod_{j=2}^n (x_j - y_j)^{2\lambda_j} dy \\ &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad - \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j (x_j - y_j))^{-\lambda_j - k_j} \frac{\partial}{\partial y_1} [(x_1 - y_1)^{\lambda_1 - k_1} f(x - y)] h_{k_1, \dots, k_n}(y) \prod_{j=2}^n (x_j - y_j)^{2\lambda_j} dy \\ &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad - \int_{\mathbb{R}^{n-1}} x_1^{-\lambda_1 - k_1} \prod_{j=2}^n \frac{(x_j - y_j)^{2\lambda_j}}{(x_j (x_j - y_j))^{\lambda_j + k_j}} \int_{-\infty}^{\infty} \frac{\partial}{\partial y_1} [(x_1 - y_1)^{\lambda_1 - k_1} f(x - y)] h_{k_1, \dots, k_n}(y) dy_1 \dots dy_n \\ &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad + \int_{\mathbb{R}^{n-1}} \lim_{\varepsilon \rightarrow 0} \{ -(x_1 - y_1)^{\lambda_1 - k_1} f(x - y) h_{k_1, \dots, k_n}(y) \}_{x_1 + \varepsilon}^{\infty} - (x_1 - y_1)^{\lambda_1 - k_1} f(x - y) h_{k_1, \dots, k_n}(y) \}_{-\infty}^{x_1 - \varepsilon} \\ &\quad + \int_{\mathbb{R} \setminus (x_1 - \varepsilon, x_1 + \varepsilon)} (x_1 - y_1)^{\lambda_1 - k_1} f(x - y) \frac{\partial}{\partial y_1} h_{k_1, \dots, k_n}(y) dy_1 \} x_1^{-\lambda_1 - k_1} \prod_{j=2}^n \frac{(x_j - y_j)^{2\lambda_j}}{(x_j (x_j - y_j))^{\lambda_j + k_j}} dy_2 \dots dy_n \\ &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad + \int_{\mathbb{R}^n} \prod_{j=1}^n (x_j (x_j - y_j))^{-\lambda_j - k_j} \prod_{j=1}^n (x_j - y_j)^{2\lambda_j} f(x - y) \frac{\partial}{\partial y_1} h_{k_1, \dots, k_n}(y) dy \\ &= -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x - y) \prod_{j=1}^n y_j^{2\lambda_j} dy \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j - k_j} f(y) \frac{\partial}{\partial u_1} h_{k_1, \dots, k_n}(u)|_{u=x-y} \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& = -(\lambda_1 + k_1) \int_{\mathbb{R}^n} x_1^{-\lambda_1 - k_1 - 1} y_1^{-\lambda_1 - k_1} \prod_{j=2}^n (x_j y_j)^{-\lambda_j - k_j} f(y) h_{k_1, \dots, k_n}(x-y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& \quad + \int_{\mathbb{R}^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j - k_j} f(y) \frac{\partial}{\partial x_1} h_{k_1, \dots, k_n}(x-y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& = \int_{\mathbb{R}^n} \frac{\partial}{\partial x_1} \left[\prod_{j=1}^n (x_j y_j)^{-\lambda_j - k_j} h_{k_1, \dots, k_n}(x-y) \right] f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

Hence, we obtain, for each $x \in (0, \infty)^n$,

$$\begin{aligned}
(41) \quad & \frac{\partial}{\partial x_1} \left(\int_{\mathbb{R}^n} f(y) \int_0^a \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \sum_{\substack{k_1, \dots, k_n \in \mathbb{N} \\ k_1, \dots, k_n \leq n \\ (k_1, \dots, k_n) \neq 0}} c_{k_1, \dots, k_n} \frac{t^{k_1 + \dots + k_n}}{\prod_{j=1}^n (x_j y_j)^{k_j}} dt \prod_{j=1}^n y_j^{2\lambda_j} dy \right) \\
& = \int_{\mathbb{R}^n} f(y) \int_0^a \frac{\partial}{\partial x_1} \left(\prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \sum_{\substack{k_1, \dots, k_n \in \mathbb{N} \\ k_1, \dots, k_n \leq n \\ (k_1, \dots, k_n) \neq 0}} c_{k_1, \dots, k_n} \frac{t^{k_1 + \dots + k_n}}{\prod_{j=1}^n (x_j y_j)^{k_j}} \right) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

Also, by using (39) we can see that

$$\begin{aligned}
(42) \quad & \frac{\partial}{\partial x_1} \int_{\mathbb{R}^n} f(y) \int_0^a \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} g_n(x, y, t) dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& = \int_{\mathbb{R}^n} f(y) \int_0^a \frac{\partial}{\partial x_1} \left(\prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} g_n(x, y, t) \right) dt \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

From (35), (41) and (42) we deduce that

$$\begin{aligned}
(43) \quad & \frac{\partial}{\partial x_1} \int_{\mathbb{R}^n} f(y) \int_0^a \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \right) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& = \int_{\mathbb{R}^n} f(y) \int_0^a \frac{\partial}{\partial x_1} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \right) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

By (32) it follows that

$$\begin{aligned}
& \left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \right| \frac{1}{\sqrt{t}} \\
& \leq \sum_{i=1}^n \frac{1}{\sqrt{t}} \prod_{j=1}^{i-1} W_t^{\lambda_j}(x_j, y_j) \left| W_t^{\lambda_i}(x_i, y_i) - \frac{t^{-\lambda_i - 1/2}}{2^{2\lambda_i} \Gamma(\lambda_i + 1/2)} \right| \frac{t^{-\sum_{j=i+1}^n (\lambda_j + 1/2)}}{\prod_{j=i+1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} + \frac{C}{t^{(n+1)/2}} \\
& \leq C \left(\frac{1}{t^{\sum_{j=1}^n (\lambda_j + 1/2) + 3/2}} + \frac{1}{t^{(n+1)/2}} \right), \quad b < t < \infty, x \in \Omega \text{ and } y \in K.
\end{aligned}$$

Moreover, according to (5) we have that

$$\begin{aligned} & \frac{\partial}{\partial x_1} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \prod_{j=1}^n \frac{(x_j y_j)^{-\lambda_j}}{2\sqrt{\pi t}} e^{-\frac{(x_j - y_j)^2}{4t}} \right) \\ &= \frac{\prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j)}{(2t)^{\lambda_1 + 1/2}} \left[\left(\frac{x_1 y_1}{2t} \right)^{-\lambda_1 + 1/2} I_{\lambda_1 + 1/2} \left(\frac{x_1 y_1}{2t} \right) \frac{y_1}{2t} - \frac{x_1}{2t} \left(\frac{x_1 y_1}{2t} \right)^{-\lambda_1 + 1/2} I_{\lambda_1 - 1/2} \left(\frac{x_1 y_1}{2t} \right) \right] e^{-\frac{x_1^2 + y_1^2}{4t}} \\ &+ \lambda_1 \frac{x_1^{-\lambda_1 - 1} y_1^{-\lambda_1}}{2\sqrt{\pi t}} \prod_{j=2}^n (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{2\sqrt{\pi t}} + \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \frac{(x_1 - y_1)}{2t}, \quad t > 0, x, y \in (0, \infty)^n. \end{aligned}$$

Then, (3) leads to

$$\begin{aligned} & \left| \frac{\partial}{\partial x_1} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \right) \right| \\ & \leq C \left(\frac{1}{t^{\sum_{j=1}^n (\lambda_j + 1/2) + 1}} + \frac{1}{t^{n/2}} \right), \quad t > b, x \in \Omega \text{ and } y \in K. \end{aligned}$$

Hence, for each $x \in (0, \infty)^n$, we can differentiate under the integral sign obtaining

(44)

$$\begin{aligned} & \frac{\partial}{\partial x_1} \int_{(0, \infty)^n} f(y) \int_b^\infty \left(W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \right) \frac{dt}{\sqrt{t}} dy \\ &= \int_{(0, \infty)^n} f(y) \prod_{j=1}^n y_j^{2\lambda_j} \int_b^\infty \frac{\partial}{\partial x_1} \left(W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \right) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{2\lambda_j} dy. \end{aligned}$$

Finally, it is not hard to see that

(45)

$$\begin{aligned} & \frac{\partial}{\partial x_1} \int_{(0, \infty)^n} f(y) \left(\int_a^b \left(W_t^{\lambda_j}(x_j, y_j) - \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \right) \frac{dt}{\sqrt{t}} - \int_1^b \frac{t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \frac{dt}{\sqrt{t}} \right) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &= \int_{(0, \infty)^n} f(y) \int_a^b \frac{\partial}{\partial x_1} \left(W_t^{\lambda_j}(x_j, y_j) - \frac{e^{-\frac{|x - y|^2}{4t}}}{(2\sqrt{\pi t})^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \right) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n. \end{aligned}$$

By combining (34), (43), (44) and (45) we can prove (33). Moreover the estimations that we have established show the absolute convergence of the integral in (33).

Thus the proof of Proposition 4.1 is finished. $\square$

Proposition 4.1 allows us to define the Riesz transforms $R_i^{\lambda_1, \dots, \lambda_n}$, $i = 1, \dots, n$, on $C_c^\infty((0, \infty)^n)$.

Proposition 4.2. *Let $f \in C_c^\infty((0, \infty)^n)$ and $\lambda_j > -1/2$, $j = 1, \dots, n$. Then, for every $i = 1, \dots, n$, the function $\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f$ admits derivative $\frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f$ with respect to x_i on almost all $(0, \infty)^n$, and*

$$\frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x - y| > \varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad \text{a.e. } x \in (0, \infty)^n.$$

Proof. Let $i = 1, \dots, n$. As it is well known, for every $g \in C_c^\infty(\mathbb{R}^n)$, $\Delta^{-1/2}g$ admits derivative $\frac{\partial}{\partial x_i} \Delta^{-1/2}g$ with respect to x_i on almost all $\mathbb{R}^n$ and

$$\frac{\partial}{\partial x_i} \Delta^{-1/2}g(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} g(y) \frac{\partial}{\partial x_i} \int_0^\infty \frac{e^{-\frac{|x-y|^2}{4t}}}{2^n(\pi t)^{(n+1)/2}} dt dy, \text{ a.e. } x \in (0, \infty)^n.$$

Moreover, for every $x \in (0, \infty)^n$, it has

$$\int_{(0, \infty)^n} |f(y)| \prod_{j=1}^n y_j^{\lambda_j} \int_0^\infty \frac{e^{-\frac{|x-y|^2}{4t}}}{t^{\frac{n+1}{2}}} dt dy \leq C \int_{(0, \infty)^n} \frac{|f(y)| \prod_{j=1}^n y_j^{\lambda_j}}{|x-y|^{n-1}} dy < \infty.$$

Then,

$$\begin{aligned} & \frac{\partial}{\partial x_i} \left(\prod_{j=1}^n x_j^{-\lambda_j} \Delta^{-1/2} \left(\prod_{j=1}^n y_j^{\lambda_j} f \right)(x) \right) \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} f(y) \frac{\partial}{\partial x_i} \left(\prod_{j=1}^n x_j^{-\lambda_j} \int_0^\infty \frac{e^{-\frac{|x-y|^2}{4t}}}{2^n(\pi t)^{(n+1)/2}} dt \right) \prod_{j=1}^n y_j^{\lambda_j} dy, \text{ a.e. } x \in (0, \infty)^n. \end{aligned}$$

Hence, from Proposition 4.1 we conclude that $\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f$ admits derivative $\frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2}$ with respect to x_i on almost all $(0, \infty)^n$ and

$$\begin{aligned} \frac{\partial}{\partial x_i} \Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) &= \frac{\partial}{\partial x_i} \left(\Delta_{\lambda_1, \dots, \lambda_n}^{-1/2} f(x) - \prod_{j=1}^n x_j^{\lambda_j} \Delta^{-1/2} \left(f \prod_{j=1}^n y_j^{\lambda_j} \right)(x) \right) \\ &\quad + \frac{\partial}{\partial x_i} \left(\prod_{j=1}^n x_j^{\lambda_j} \Delta^{-1/2} \left(f \prod_{j=1}^n y_j^{\lambda_j} \right)(x) \right) \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} f(y) \left(R_i^{\lambda_1, \dots, \lambda_n}(x, y) - \mathcal{R}_i^{\lambda_1, \dots, \lambda_n}(x, y) \right) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\quad + \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} f(y) \mathcal{R}_i^{\lambda_1, \dots, \lambda_n}(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} f(y) R_i^{\lambda_1, \dots, \lambda_n}(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \text{ a.e. } x \in (0, \infty)^n. \end{aligned}$$

□

We now prove that, for every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, exists the limit

$$\lim_{\varepsilon \rightarrow 0^+} \int_{|x-y|>\varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \text{ a.e. } x \in (0, \infty)^n,$$

with $i = 1, \dots, n$. In order to show this we consider, for every $i = 1, \dots, n$, the maximal operator $R_{i,*}^{\lambda_1, \dots, \lambda_n}$ defined by

$$R_{i,*}^{\lambda_1, \dots, \lambda_n}(f)(x) = \sup_{\varepsilon > 0} \left| \int_{|x-y|>\varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|.$$

Proposition 4.3. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$, and $i = 1, \dots, n$. The maximal operator $R_{i,*}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, $1 < p < \infty$ and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

Proof. We consider $i = 1$. For the other values of i we can proceed analogously. We can write

$$\begin{aligned}
R_1^{\lambda_1, \dots, \lambda_n}(x, y) &= \frac{1}{\sqrt{\pi}} \int_0^\infty \left(\frac{\partial}{\partial x_1} W_t^{\lambda_1}(x_1, y_1) - \chi_{\{x_1/2 < y_1 < 2x_1\}}(y_1)(x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \right) \\
&\times \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) \frac{dt}{\sqrt{t}} \\
&+ \sum_{l=2}^n \frac{1}{\sqrt{\pi}} \int_0^\infty (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \prod_{j=2}^{l-1} (x_j y_j)^{-\lambda_j} \mathbb{W}_t(x_j, y_j) \\
&\times \prod_{j=1}^{l-1} \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j) \left(W_t^{\lambda_l}(x_l, y_l) - \chi_{\{x_l/2 < y_l < 2x_l\}}(y_l)(x_l y_l)^{-\lambda_l} \mathbb{W}_t(x_l, y_l) \right) \prod_{j=l+1}^n W_t^{\lambda_j}(x_j, y_j) \frac{dt}{\sqrt{t}} \\
&+ \frac{1}{\sqrt{\pi}} \int_0^\infty (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \prod_{j=2}^n (x_j y_j)^{-\lambda_j} \mathbb{W}_t(x_j, y_j) \prod_{j=1}^n \chi_{\{x_j/2 < y_j < 2x_j\}}(y_j) \frac{dt}{\sqrt{t}} \\
&= \sum_{l=1}^{n+1} R_{1,l}^{\lambda_1, \dots, \lambda_n}(x, y), \quad x, y \in (0, \infty)^n.
\end{aligned}$$

We now study the maximal operator $R_{1,l,*}^{\lambda_1, \dots, \lambda_n}$ associated with $R_{1,l}^{\lambda_1, \dots, \lambda_n}$ in the usual way. Firstly we consider $R_{1,1,*}^{\lambda_1, \dots, \lambda_n}$. According to [22, (22)], (3) and (4), we have

$$(46) \quad \left| \frac{\partial}{\partial u} W_t^\lambda(u, v) \right| \leq C \frac{u+v}{t^{\lambda+3/2}} e^{-(u^2+v^2)/4}, \quad t, u, v \in (0, \infty) \text{ and } \frac{uv}{t} \leq 1,$$

and

$$(47) \quad \left| \frac{\partial}{\partial u} W_t^\lambda(u, v) \right| \leq C(uv)^{-\lambda} \frac{e^{-(u-v)^2/8t}}{t}, \quad t, u, v \in (0, \infty) \text{ and } \frac{uv}{t} \geq 1,$$

Inequalities (46) and (47) lead, for every $\lambda > -1/2$ and $0 < v < u/2$, to

$$(48) \quad \left| \frac{\partial}{\partial u} W_t^\lambda(u, v) \right| \leq C \frac{ue^{-u^2/32t}}{t^{\lambda+3/2}} \leq C \frac{e^{-u^2/40t}}{t^{\lambda+1}}, \quad t > 0$$

Suppose that $1 \leq l \leq m \leq n$. According to (6), (9), (12) and (48), we get

$$\begin{aligned}
&\int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_l}{2}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \dots \int_{\frac{x_m}{2}}^{2x_m} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty |R_{1,1}^{\lambda_1, \dots, \lambda_n}(x, y)| |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_l}{2}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \dots \int_{\frac{x_m}{2}}^{2x_m} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty \prod_{j=m+1}^n \frac{1}{y_j^{2\lambda_j+1}} |f(y)| \\
&\quad \times \int_0^\infty \frac{e^{-(\sum_{j=1}^l x_j^2)/40t}}{t^{\sum_{j=1}^l (\lambda_j+1/2)+1}} \prod_{j=l+1}^m \left(\frac{1}{x_j^{2\lambda_j+1}} + \frac{(x_j y_j)^{-\lambda_j}}{\sqrt{t}} e^{-(x_j-y_j)^2/4t} \right) dt \prod_{j=1}^n y_j^{2\lambda_j} dy
\end{aligned}$$

Since, for every $k \in \mathbb{N}$, H_∞^k and $H_{loc}^{\alpha_1, \dots, \alpha_k}$ are bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself, provided that $\alpha_j > -1/2$, $j = 1, \dots, k$, the operator

$$S_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) = \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_l}{2}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \dots \int_{\frac{x_m}{2}}^{2x_m} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty |R_{1,1}^{\lambda_1, \dots, \lambda_n}(x, y)| |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, provided that, for every $r, k \in \mathbb{N}$, $1 < r \leq k$ and $\alpha_j > -1/2$, $j = 1, \dots, k$, the operator

$$\begin{aligned} \mathcal{S}_r^{\alpha_1, \dots, \alpha_k}(g)(x) &= \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_r}{2}} \int_{\frac{x_{r+1}}{2}}^{2x_{r+1}} \cdots \int_{\frac{x_k}{2}}^{2x_k} |g(y)| \\ &\quad \times \int_0^\infty \prod_{j=r+1}^k (x_j y_j)^{-\alpha_j} \frac{e^{-(\sum_{j=1}^r x_j^2 + \sum_{j=r+1}^k (x_j - y_j)^2)/40t}}{\sum_{j=1}^r (\alpha_j + 1/2) + 1 + (k-r)/2} dt \prod_{j=1}^k y_j^{2\alpha_j} dy \\ &= c_{\alpha_1, \dots, \alpha_k}^{k,r} \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_r}{2}} \int_{\frac{x_{r+1}}{2}}^{2x_{r+1}} \cdots \int_{\frac{x_k}{2}}^{2x_k} \prod_{j=r+1}^k (x_j y_j)^{-\alpha_j} |g(y)| \\ &\quad \times \frac{1}{\left(\sum_{j=1}^r x_j^2 + \sum_{j=r+1}^k (x_j - y_j)^2\right)^{\sum_{j=1}^r (\alpha_j + 1/2) + (k-r)/2}} \prod_{j=1}^k y_j^{2\alpha_j} dy, \end{aligned}$$

where $c_{\alpha_1, \dots, \alpha_k}^{k,r}$ is a constant, is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$. This last property is true because the operators $\mathcal{H}_{r,k}^{\alpha_1, \dots, \alpha_k}$ and $L_{\alpha_1, \dots, \alpha_k}$ are bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, when $\alpha_j > -1/2$, $j = 1, \dots, k$, and $1 < r \leq k$, $r, k \in \mathbb{N}$.

On the other hand, as it was proved in [8, Lemma 5],

$$\int_0^\infty \left| \frac{\partial}{\partial x_1} W_t^{\lambda_1}(x_1, y_1) \right| \frac{dt}{\sqrt{t}} \leq C \frac{1}{x_1^{2\lambda_1+1}}, \quad 0 < y_1 < x_1/2.$$

Then, by writing $\bar{y} = (y_2, \dots, y_n)$, it has, for every $x \in (0, \infty)^n$,

$$\begin{aligned} S_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) &\leq C \int_0^{\frac{x_1}{2}} \int_0^\infty \left| \frac{\partial}{\partial x_1} W_t^{\lambda_1}(x_1, y_1) \right| \\ &\quad \times \left(\sup_{t>0} \int_{(0,\infty)^{n-1}} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y_1, \bar{y})| \prod_{j=2}^n y_j^{2\lambda_j} d\bar{y} \right) dt y_1^{2\lambda_1} dy_1 \\ &\leq \frac{C}{x_1^{2\lambda_1+1}} \int_0^{\frac{x_1}{2}} \sup_{t>0} \int_{(0,\infty)^{n-1}} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y_1, \bar{y})| \prod_{j=2}^n y_j^{2\lambda_j} d\bar{y} y_1^{2\lambda_1} dy_1. \end{aligned}$$

Since L_{λ_1} is bounded from $L^p((0, \infty), x^{2\lambda_1} dx)$ into itself and $W_*^{\lambda_2, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=2}^n x_j^{2\alpha_j} d\bar{x})$ into itself, $S_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\alpha_j} dx)$ into itself, for every $1 < p < \infty$.

By using (46) and (47), for every $\lambda > -1/2$ and $2u < v < \infty$, we obtain

$$(49) \quad \left| \frac{\partial}{\partial u} W_t^\lambda(u, v) \right| \leq \frac{C}{t^{\lambda+1}} e^{-v^2/40t}, \quad t > 0.$$

Suppose that $1 \leq l \leq m \leq n$. By (6), (9), (12) and (49), for each $x = (x_1, \dots, x_n) \in (0, \infty)^n$, it follows that,

$$\mathcal{S}_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) = \int_{2x_1}^\infty \cdots \int_{2x_l}^\infty \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \cdots \int_{\frac{x_m}{2}}^{2x_m} \int_0^{\frac{x_{m+1}}{2}} \cdots \int_0^{\frac{x_n}{2}} |R_{1,1}^{\lambda_1, \dots, \lambda_n}(x, y)| |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy$$

$$\begin{aligned} &\leq C \int_{2x_1}^{\infty} \cdots \int_{2x_l}^{\infty} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \cdots \int_{\frac{x_m}{2}}^{2x_m} \int_0^{\frac{x_{m+1}}{2}} \cdots \int_0^{\frac{x_n}{2}} |f(y)| \int_0^{\infty} \frac{e^{-(\sum_{j=1}^l y_j^2 + \sum_{j=m+1}^n x_j^2)/40t}}{t^{\sum_{j=1}^l (\lambda_j+1/2) + \sum_{j=m+1}^n (\lambda_j+1/2)+1}} \\ &\quad \times \prod_{j=l+1}^m \left[(x_j y_j)^{-\lambda_j} \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{t}} + \frac{1}{x_j^{2\lambda_j+1}} \right] dt \prod_{j=1}^n y_j^{2\lambda_j} dy. \end{aligned}$$

Since the operator $H_{loc}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into itself, provided that $\alpha_j > -1/2$, $j = 1, \dots, k$, in order that to see that the operator $\mathcal{S}_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ it is sufficient to show that, for every $\alpha_j > -1/2$, $j = 1, \dots, k$, and $1 < s \leq r \leq k$, $s, r, k \in \mathbb{N}$, the operator

$$\begin{aligned} T_{s,r}^{\alpha_1, \dots, \alpha_k}(g)(x) &= \int_{2x_1}^{\infty} \int_{2x_2}^{\infty} \cdots \int_{2x_s}^{\infty} \int_{\frac{x_{s+1}}{2}}^{2x_{s+1}} \cdots \int_{\frac{x_r}{2}}^{2x_r} \int_0^{\frac{x_{r+1}}{2}} \cdots \int_0^{\frac{x_k}{2}} |g(y)| \prod_{j=1}^k y_j^{2\alpha_j} \\ &\quad \times \frac{\prod_{j=s+1}^r (x_j y_j)^{-\alpha_j} dy}{\left(\sum_{j=1}^s y_j^2 + \sum_{j=s+1}^r (x_j - y_j)^2 + \sum_{j=r+1}^k x_j^2 \right)^{\sum_{j=1}^s (\lambda_j+1/2) + \sum_{j=r+1}^k (\lambda_j+1/2) + (r-s)/2}}, \quad x \in (0, \infty)^k, \end{aligned}$$

is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$. Let $\alpha_j > -1/2$, $j = 1, \dots, k$, and $0 < s \leq r \leq k$. By proceeding as in the proof of [25, Case 3] we can prove that the operator $T_{s,r}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, when $s < r$. Also, the operator $T_{s,s}^{\alpha_1, \dots, \alpha_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\alpha_j} dx)$, because

$$T_{s,s}(g)(x) \leq \frac{C}{\left(\sum_{j=s+1}^k x_j^2 \right)^{\sum_{j=s+1}^k (\lambda_j+1/2)}} \int_0^{\frac{x_{s+1}}{2}} \cdots \int_0^{\frac{x_{k+1}}{2}} \int_{2x_1}^{\infty} \cdots \int_{2x_s}^{\infty} \frac{|g(y)|}{y_1 \cdots y_s} \prod_{j=s+1}^k y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^k,$$

and H_{∞}^s is bounded from $L^1((0, \infty)^s, \prod_{j=1}^s x_j^{2\lambda_j} dx)$ into itself and $L_{\lambda_{s+1}, \dots, \lambda_k}$ is bounded from $L^1((0, \infty)^{k-s}, \prod_{j=s+1}^k x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^{k-s}, \prod_{j=s+1}^k x_j^{2\lambda_j} dx)$. Then, the operator $\mathcal{S}_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Moreover, [8, Lemma 5],

$$\int_0^{\infty} \left| \frac{\partial}{\partial x_1} W_t^{\lambda_1}(x_1, y_1) \right| \frac{dt}{\sqrt{t}} \leq C \frac{x_1}{y_1^{2\lambda_1+2}}, \quad 2x_1 < y_1 < \infty.$$

Then, for each $x \in (0, \infty)^n$, putting $\bar{y} = (y_2, \dots, y_n)$,

$$\mathcal{S}_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) \leq C \int_{2x_1}^{\infty} \frac{1}{y_1} \left(\sup_{t>0} \int_{(0, \infty)^{n-1}} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y_1, \bar{y})| \prod_{j=2}^n y_j^{2\lambda_j} d\bar{y} \right) dy_1.$$

Since H_{∞}^1 is bounded from $L^p((0, \infty), x^{2\lambda_1} dx)$ into itself and $W_*^{\lambda_2, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^{n-1}, \prod_{j=2}^n x_j^{2\lambda_j} dx)$ into itself, $\mathcal{S}_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

Finally, as it was proved in the proof of [8, Lemma 3], we have that

$$\left| \frac{\partial}{\partial u} W_t^\lambda(u, v) + (uv)^{-\lambda} \frac{u-v}{4\sqrt{\pi}t^{3/2}} e^{-(u-v)^2/4t} \right| \leq C \frac{e^{-(u-v)^2/4t}}{\sqrt{t}} (uv)^{-\lambda-1/2}, \quad uv \geq t.$$

Then, by using (6) and (9), we obtain that, for every $1 \leq l \leq m \leq n$,

$$\begin{aligned} \mathbb{S}_{l,m}^{\lambda_1, \dots, \lambda_n}(f)(x) &= \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_l}{2}}^{2x_l} \int_0^{\frac{x_{l+1}}{2}} \dots \int_0^{\frac{x_m}{2}} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty |R_{1,1}^{\lambda_1, \dots, \lambda_n}(x, y)| |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\leq C \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_l}{2}}^{2x_l} \int_0^{\frac{x_{l+1}}{2}} \dots \int_0^{\frac{x_m}{2}} \int_{2x_{m+1}}^\infty \dots \int_{2x_n}^\infty |f(y)| \left[\int_0^{x_1 y_1} \frac{e^{-(x_1 - y_1)^2/4t}}{t} (x_1 y_1)^{-\lambda_1 - 1/2} \right. \\ &\quad \times \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) dt + \int_{x_1 y_1}^\infty \left(\frac{e^{-x_1^2/8t}}{t^{\lambda_1+3/2}} + (x_1 y_1)^{-\lambda_1} \frac{e^{-(x_1 - y_1)^2/8t}}{t^{3/2}} \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) dt \Big] \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\leq C \left[\int_0^{2x_1^2} \frac{dt}{\sqrt{t}} \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \int_{(0,\infty)^{n-1}} \frac{e^{-(x_1 - y_1)^2/4t}}{\sqrt{t}} (x_1 y_1)^{-\lambda_1 - 1/2} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \right. \\ &\quad \left. + \left(\int_{\frac{x_1^2}{2}}^\infty \frac{dt}{t^{\lambda_1+3/2}} + x_1^{-2\lambda_1} \int_{\frac{x_1^2}{2}}^\infty \frac{dt}{t^{3/2}} \right) \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \int_{(0,\infty)^{n-1}} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \right] \\ &\leq C \left[\sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \int_{(0,\infty)^{n-1}} \frac{e^{-(x_1 - y_1)^2/4t}}{\sqrt{t}} (x_1 y_1)^{-\lambda_1} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \right. \\ &\quad \left. + \sup_{t>0} \int_{(0,\infty)^{n-1}} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) \frac{1}{x_1^{2\lambda_1+1}} \int_{\frac{x_1}{2}}^{2x_1} f(y) y_1^{2\lambda_1} dy_1 \prod_{j=2}^n y_j^{2\lambda_j} dy \right]. \end{aligned}$$

Since the operator $H_{loc}^{\lambda_1}$ is bounded from $L^1((0, \infty), x_1^{2\lambda_1+1} dx_1)$ into itself, the maximal operator $W_*^{\lambda_2, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^{n-1}, \prod_{j=2}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^{n-1}, \prod_{j=2}^n x_j^{2\lambda_j} dx)$ (Theorem 1.1), and, as the arguments developed in the proof of Theorem 1.1 show, the maximal operator

$$\mathcal{W}_*^{\lambda_1, \dots, \lambda_n}(f)(x) = \sup_{t>0} \int_{\frac{x_1}{2}}^{2x_1} \int_{(0,\infty)^{n-1}} \frac{e^{-(x_1 - y_1)^2/4t}}{\sqrt{t}} (x_1 y_1)^{-\lambda_1} \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, for each $x \in (0, \infty)^n$. We conclude that the operator $\mathbb{S}_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Moreover, since the operator $H_{loc}^{\lambda_1}$, $W_*^{\lambda_2, \dots, \lambda_n}$ and $W_*^{\lambda_1, \dots, \lambda_n}$ are bounded from $L^p((0, \infty), x^{2\lambda_1} dx)$ into itself, $L^p((0, \infty)^{n-1}, \prod_{j=2}^n x_j^{2\lambda_j} dx)$ into itself, and $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, respectively, $\mathbb{S}_{l,m}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

Hence, we have proved that the operator $R_{1,1}^{\lambda_1, \dots, \lambda_n}$ defined by

$$R_{1,1}^{\lambda_1, \dots, \lambda_n}(f)(x) = \int_{(0,\infty)^n} |R_{1,1}^{\lambda_1, \dots, \lambda_n}(x, y)| f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n,$$

is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ and from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$. Since

$$\begin{aligned} R_{1,1,*}^{\lambda_1,\dots,\lambda_n}(f)(x) &= \sup_{\varepsilon>0} \left| \int_{|x-y|>\varepsilon} R_{1,1}^{\lambda_1,\dots,\lambda_n}(x,y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| \\ &\leq R_{1,1}^{\lambda_1,\dots,\lambda_n}(f)(x), \quad x \in (0, \infty)^n, \end{aligned}$$

$R_{1,1,*}^{\lambda_1,\dots,\lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ and from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

In order to study the maximal operators $R_{1,l,*}^{\lambda_1,\dots,\lambda_n}$, $l = 2, \dots, n$, we can proceed as for $R_{1,1,*}^{\lambda_1,\dots,\lambda_n}$ by taking into account (29).

We now consider the maximal operator

$$R_{1,n+1,*}^{\lambda_1,\dots,\lambda_n}(f)(x) = \sup_{\varepsilon>0} \left| \int_{|x-y|>\varepsilon} R_{1,n+1}^{\lambda_1,\dots,\lambda_n}(x,y) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|, \quad x \in (0, \infty)^n.$$

We use now a procedure similar than the one developed in the proof of Claim 1 in Section 3.

For every $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$ we define the operator $\mathcal{Y}_m^{\lambda_1,\dots,\lambda_n}$ by

$$\begin{aligned} \mathcal{Y}_m^{\lambda_1,\dots,\lambda_n}(f)(x) &= \sup_{\varepsilon>0} \left| \chi_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})}(x) \prod_{j=1}^n x_j^{-\lambda_j} \frac{1}{\sqrt{\pi}} \int_{|x-y|>\varepsilon} f_m(y) \cdot \right. \\ &\quad \cdot \left. \int_0^\infty \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \prod_{j=2}^n \mathbb{W}_t(x_j, y_j) \frac{dt}{\sqrt{t}} dy \right|, \quad x \in (0, \infty)^n, \end{aligned}$$

where $f_m(y) = \chi_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})}(y) f(y) \prod_{j=1}^n y_j^{\lambda_j}$, $y \in (0, \infty)^n$.

Let $\gamma > 0$. We have that

$$\begin{aligned} &m_{\lambda_1,\dots,\lambda_n}(\{x \in (0, \infty)^n : R_{1,n+1,*}^{\lambda_1,\dots,\lambda_n}(f)(x) > \gamma\}) \\ &\leq m_{\lambda_1,\dots,\lambda_n}(\{x \in (0, \infty)^n : |R_{1,n+1,*}^{\lambda_1,\dots,\lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} \mathcal{Y}_m^{\lambda_1,\dots,\lambda_n}(f)(x)| > \gamma/2\}) \\ &\quad + \sum_{m \in \mathbb{Z}^n} m_{\lambda_1,\dots,\lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \mathcal{Y}_m^{\lambda_1,\dots,\lambda_n}(f)(x) > \gamma/2\}). \end{aligned}$$

We denote by $R_{1,*}$ the maximal operator

$$R_{1,*}(g)(x) = \sup_{\varepsilon>0} \left| \int_{|x-y|>\varepsilon} \int_0^\infty \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \prod_{j=2}^n \mathbb{W}_t(x_j, y_j) \frac{dt}{\sqrt{t}} g(y) dy \right|, \quad x \in \mathbb{R}^n.$$

It is well known that $R_{1,*}$ is a bounded operator from $L^1(\mathbb{R}^n, dx)$ into $L^{1,\infty}(\mathbb{R}^n, dx)$ and from $L^p(\mathbb{R}^n, dx)$ into itself, for every $1 < p < \infty$.

Then, for every $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$, it gets

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}) \\
& \leq C 2^{\sum_{j=1}^n \lambda_j m_j} m_0^{(n)}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : R_{1,*}(f_m)(x) > M\gamma 2^{\sum_{j=1}^n \lambda_j m_j}\}) \\
& \leq \frac{C}{\gamma} 2^{\sum_{j=1}^n \lambda_j m_j} \|f_m\|_{L^1(\mathbb{R}^n, dx)} \\
& \leq \frac{C}{\gamma} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

Here M and C denotes suitable positive constants that do not depend on $m \in \mathbb{Z}^n$. By $m_0^{(n)}$ we represent the Lebesgue measure in $\mathbb{R}^n$. Hence,

$$\begin{aligned}
(50) \quad & \sum_{m \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x) > \gamma/2\}) \\
& \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j})}.
\end{aligned}$$

Also, if $1 < p < \infty$, we can write

$$\begin{aligned}
(51) \quad & \left\| \sum_{m \in \mathbb{Z}^n} \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f) \right\|_p^p = \sum_{m \in \mathbb{Z}^n} \int_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})} |\mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)|^p \prod_{j=1}^n x_j^{2\lambda_j} dx \\
& \leq C \sum_{m \in \mathbb{Z}^n} 2^{(2-p)\sum_{j=1}^n m_j \lambda_j} \int_{\mathbb{R}^n} |R_{1,*}(f_m)(x)|^p dx \\
& \leq C \sum_{m \in \mathbb{Z}^n} 2^{(2-p)\sum_{j=1}^n m_j \lambda_j} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)|^p \prod_{j=1}^n y_j^{p\lambda_j} dy \\
& \leq C \sum_{m \in \mathbb{Z}^n} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)|^p \prod_{j=1}^n y_j^{2\lambda_j} dy \\
& \leq C \|f\|_p^p.
\end{aligned}$$

On the other hand, we have that

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |R_{1,n+1,*}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \\
& \leq \sum_{m \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}(\{x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : |R_{1,n+1,*}^{\lambda_1, \dots, \lambda_n}(f)(x) - \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}).
\end{aligned}$$

Let $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$. We define, for every $x \in \prod_{j \in \mathbb{N}} [2^{m_j}, 2^{m_j+1})$, the sets $B_l(x)$ and $C_l(x)$, $l = 1, \dots, n$, and $A(x)$ as in the proof of Theorem 1.2. Then,

$$\begin{aligned}
& |R_{1,n+1,*}^{\lambda_1, \dots, \lambda_n}(f)(x) - \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)| \\
& \leq \frac{1}{\sqrt{\pi}} \sum_{l=1}^n \left(\prod_{j=1}^n x_j^{-\lambda_j} \int_{B_l(x)} |f(y)| \int_0^\infty \left| \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \right| \prod_{j=2}^n \mathbb{W}_t(x_2, y_2) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{\lambda_j} dy \right. \\
& \quad \left. + \prod_{j=1}^n x_j^{-\lambda_j} \int_{C_l(x)} |f(y)| \int_0^\infty \left| \frac{\partial}{\partial x_1} \mathbb{W}_t(x_1, y_1) \right| \prod_{j=2}^n \mathbb{W}_t(x_2, y_2) \frac{dt}{\sqrt{t}} \prod_{j=1}^n y_j^{\lambda_j} dy \right)
\end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{l=1}^n \left(\int_{B_l(x)} |f(y)| \int_0^\infty \frac{e^{-|x-y|^2/8t}}{t^{n/2+1}} dt dy + \int_{C_l(x)} |f(y)| \int_0^\infty \frac{e^{-|x-y|^2/8t}}{t^{n/2+1}} dt dy \right) \\
&\leq C \sum_{l=1}^n \left(\int_{B_l(x)} \frac{|f(y)|}{|x-y|^n} dy + \int_{C_l(x)} \frac{|f(y)|}{|x-y|^n} dy \right), \quad x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}).
\end{aligned}$$

By denoting $\bar{y} = (y_2, \dots, y_n) \in \mathbb{R}^{n-1}$, and $\tilde{f}_m(\bar{y}) = \chi_{\prod_{j=2}^n [2^{m_j-1}, 2^{m_j+2})}(\bar{y}) \int_{2^{m_1-1}}^{2^{m_1+2}} |f(y_1, \bar{y})| dy_1$, we obtain

$$\int_{B_1(x)} \frac{|f(y)|}{|x-y|^n} dy \leq C \frac{1}{2^{m_1}} M_{n-1}(\tilde{f}_m)(\bar{x}),$$

where $\bar{x} = (x_2, \dots, x_n)$ when $x = (x_1, x_2, \dots, x_n)$, and $M_{n-1}(\tilde{f}_m)(\bar{x})$ denotes the Hardy Littlewood maximal function in $\mathbb{R}^{n-1}$.

Then, for every $\beta > 0$,

$$\begin{aligned}
&m_{\lambda_1, \dots, \lambda_n} \left(\left\{ x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \int_{B_1(x)} \frac{f(y)}{|x-y|^n} dy > \beta \right\} \right) \\
&\leq m_{\lambda_1, \dots, \lambda_n} \left(\left\{ x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : \frac{C}{2^{m_1}} M_{n-1}(\tilde{f}_m)(\bar{x}) > \beta \right\} \right) \\
&\leq M 2^{m_1} 2^{2 \sum_{j=1}^n \lambda_j m_j} m_0^{(n-1)} \left(\left\{ x \in \mathbb{R}^{n-1} : M_{n-1}(\tilde{f}_m)(\bar{x}) > \frac{\beta 2^{m_1}}{C} \right\} \right) \\
(52) \quad &\leq \frac{M}{\beta} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy,
\end{aligned}$$

where $M > 0$ does not depend on $m \in \mathbb{Z}^n$. Also, if $1 < p < \infty$, by using Jensen's inequality and maximal Theorem we have

$$\begin{aligned}
&\int_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})} \left| \int_{B_1(x)} \frac{|f(y)|}{|x-y|^n} \right|^p \prod_{j=1}^n x_j^{2\lambda_j} dx \leq C 2^{2 \sum_{j=1}^n \lambda_j m_j - m_1(p-1)} \int_{\mathbb{R}^{n-1}} |M_{n-1}(\tilde{f}_m)(\bar{x})|^p dx \\
(53) \quad &\leq C \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)|^p \prod_{j=1}^n y_j^{2\lambda_j} dy,
\end{aligned}$$

where $C > 0$ does not depend on $m \in \mathbb{Z}^n$.

Estimates similar than (52) and (53) can be obtain for $B_l(x)$, $l = 2, \dots, n$, and $C_l(x)$, $l = 1, \dots, n$. Putting together all the estimates we get

$$\begin{aligned}
&m_{\lambda_1, \dots, \lambda_n} \left(\left\{ x \in \prod_{j=1}^n [2^{m_j}, 2^{m_j+1}) : |R_{1,n+1,*}^{\lambda_1, \dots, \lambda_n}(f)(x) - \mathcal{D}_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2 \right\} \right) \\
&\leq \frac{C}{\gamma} \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+2})} |f(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy
\end{aligned}$$

and, for every $1 < p < \infty$,

$$\begin{aligned}
&\int_{\prod_{j=1}^n [2^{m_j}, 2^{m_j+1})} |R_{1,n+1,*}^{\lambda_1, \dots, \lambda_n}(f)(x) - \mathcal{D}_m^{\lambda_1, \dots, \lambda_n}(f)(x)|^p \prod_{j=1}^n x_j^{2\lambda_j} dx \\
&\leq C \int_{\prod_{j=1}^n [2^{m_j-1}, 2^{m_j+1})} |f(y)|^p \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

Hence, by summing in $m \in \mathbb{Z}^n$, it follows

$$(54) \quad m_{\lambda_1, \dots, \lambda_n}(\{x \in (0, \infty)^n : |R_{1, n+1, *}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)| > \gamma/2\}) \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j})},$$

and, for every $1 < p < \infty$,

$$(55) \quad \|R_{1, n+1, *}^{\lambda_1, \dots, \lambda_n}(f)(x) - \sum_{m \in \mathbb{Z}^n} \mathcal{Y}_m^{\lambda_1, \dots, \lambda_n}(f)(x)\|_p \leq C \|f\|_p.$$

From (50) and (54) we deduce that $R_{1, n+1, *}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, and from (51) and (55) it infers that $R_{1, n+1, *}^{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

Thus the proof of Proposition 4.3 is finished. $\square$

According to Proposition 4.2 and 4.3 standard arguments allow us to conclude that, for every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, and $i = 1, \dots, n$, there exists the limit

$$\lim_{\varepsilon \rightarrow 0^+} \int_{|x-y| > \varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) dy, \quad \text{a. e. } x \in (0, \infty)^n.$$

We define, for every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, and $i = 1, \dots, n$, the Riesz transform $R_i^{\lambda_1, \dots, \lambda_n}(f)$ of f by

$$R_i^{\lambda_1, \dots, \lambda_n}(f)(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{|x-y| > \varepsilon} R_i^{\lambda_1, \dots, \lambda_n}(x, y) f(y) dy, \quad \text{a.e. } x \in (0, \infty)^n.$$

Note that, by Proposition 4.2, for every $i = 1, \dots, n$ this definition extends the initial definition of Riesz transform $R_i^{\lambda_1, \dots, \lambda_n}$ from $C_c^\infty((0, \infty)^n)$ to $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$.

Finally, from Proposition 4.3 we infer the desired L^p -boundedness properties for the Riesz transform $R_i^{\lambda_1, \dots, \lambda_n}$, $i = 1, \dots, n$, and the proof of Theorem 1.3 is complete.

REFERENCES

- [1] H. Aimar, L. Forzani, and R. Scotto, *On Riesz transform and maximal functions in the context of Gaussian harmonic analysis*, Trans. Amer. Math. Soc., 359(5) (2007), 2137-2154.
- [2] K.F. Andersen, *Norm inequalities for ultraspherical and Hankel conjugate functions*, Canad. J. Math., 27 (1975), 162-171.
- [3] K.F. Andersen, and R.A. Kerman, *Weighted norm inequalities for generalized Hankel conjugate transformations*, Studia Math., 71 (1981), 15-26.
- [4] J.J. Betancor, J.C. Fariña, M.T. Martínez, and L. Rodríguez-Mesa, *Higher order Riesz transforms associated with Bessel operators*, Arkiv Mat. 46(2) (2008), 219-250.
- [5] J.J. Betancor, J.C. Fariña, M.T. Martínez, and J.L. Torrea, *Riesz transforms and g-functions associated with Bessel operators and their appropriate Banach spaces*, Israel J. Math., 141 (2004) 125-144.
- [6] J.J. Betancor, J.C. Fariña, M.T. Martínez, and J.L. Torrea, *Riesz transforms related to Bessel operators*, Proc. Roy. Soc. Edinburgh, Section A, 135 (2005), 85-104.
- [7] J.J. Betancor, J.C. Fariña, and A. Sanabria, *On Littlewood-Paley functions associated with Bessel operators*, Glasgow Math. J., 51(1) (2009) 55-70.
- [8] J.J. Betancor, E. Harboure, A. Nowak, and B. Viviani, *Mapping properties of fundamental operators in harmonic analysis related to Bessel operators*, to appear in Studia Mathematica (available in arXiv:0802.3497).
- [9] J.J. Betancor, and K. Stempak *On Hankel conjugate functions*, Studia Sci. Math. Hungar. 41 (2004), 59-91.
- [10] L.A. Caffarelli, and C.P. Calderón, *On Abel summability of multiple Jacobi series*, Colloq. Math., 30 (1974), 277-288.
- [11] A. Chicco Ruiz, and E. Harboure, *Weighted norm inequalities for the heat diffusion Laguerre's semigroups*, Math. Z., 257 (2007), 329-354.
- [12] O. Ciaurri, A. Nowak, and K. Stempak, *Jacobi transplantation revisited*, Math. Z., 257 (2007), 355-380.

- [13] O. Ciaurri, and L. Roncal, *The Bochner-Riesz means for Fourier-Bessel expansions*, J. Funct. Anal., 228 (2005), 89-113.
- [14] U. Dinger, *Weak type $(1,1)$ estimates of the maximal function for the Laguerre semigroup in finite dimensions*, Revista matemática Iberoamericana, 8(1) (1992), 93-118.
- [15] J. García-Cuerva, G. Manceri, S. Meda, P. Sjögren, and J.L. Torrea, *Maximal operators for the holomorphic Ornstein-Uhlenbeck semigroup*, J. London Math. Soc., 67(2) (2003), 219-134.
- [16] G. Garrigós, E. Harboure, T. Signes, J.L. Torrea, and B. Viviani, *A sharp weighted transplantation theorem for Laguerre function expansions*, J. Func. Anal., 244 (2007), 247-276.
- [17] R.A. Kerman, *Generalized Hankel conjugate transformations on rearrangement invariant spaces*, Trans. Amer. Math. Soc., 232 (1977), 111-130.
- [18] R.A. Kerman, *Boundedness criteria for generalized Hankel conjugate transformations*, Canad. J. Math., 30 (1978), 147-153.
- [19] N.N. Lebedev, *Special functions and their applications*, Dover publications, New York, 1972.
- [20] R. Macías, C. Segovia, and J.L. Torrea, *Heat diffusion maximal operators for Laguerre semigroups with negative parameters*, J. Funct. Anal., 229 (2005), 300-316.
- [21] B. Muckenhoupt, *Poisson integrals for Hermite and Laguerre expansions*, Trans. Amer Math. Soc., 139 (1969), 231-242.
- [22] B. Muckenhoupt, and E. Stein, *Classical expansions and their relation to conjugate harmonic functions*, Trans. Amer. Math. Soc., 118 (1965), 17-92.
- [23] A. Nowak, and P. Sjögren, *Weak type $(1,1)$ estimates for maximal operators associated with various multi-dimensional systems of Laguerre functions*, Indiana Univ Math. J., 56 (2007), 417-436.
- [24] A. Nowak, and P. Sjögren, *The multidimensional pencil phenomenon for Laguerre heat diffusion maximal operators*, Math. Annalen, 344(1) (2009), 213-248.
- [25] A. Nowak, and K. Stempak, *Weighted estimates for the Hankel transform transplantation operator*, Tohoku Math. J., 58 (2006), 277-301.
- [26] A. Nowak, and K. Stempak, *Riesz transform for multidimensional Laguerre function expansions*, Adv. Math., 215 (2) (2007), 642-678.
- [27] E. Sasso, *Functional calculus for the Laguerre operator*, Math. Z., 249 (2005), 683-711.
- [28] P. Sjögren, *On the maximal function for the Mehler kernel*, Harmonic Analysis, Cartona, 1982, Lecture Notes in Mathematics 992 (Springer, 1983), 73-82.
- [29] E.M. Stein, *Topics in harmonic analysis related to the Littlewood-Paley theory*, Ann. of Math. Studies, vol. 63, Princeton Univ. Press, Princeton, NJ, 1970.
- [30] E.M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton University Press, Princeton, NJ, 1993.
- [31] K. Stempak, *The Littlewood-Paley theory for the Fourier-Bessel transform*, Preprint 45, Mathematical Institute, University of Wrocław, 1985.
- [32] K. Stempak, and J.L. Torrea, *Poisson integrals and Riesz transform for Hermite function expansions with weights*, J. Funct. Anal., 202 (2003), 443-472.
- [33] S. Thangavelu, *Lecture on Hermite and Laguerre expansions*, Mathematical Notes 42, Princeton University Press, Princeton, 1993.
- [34] G.N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, Cambridge, 1966.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA, CAMPUS DE ANCHIETA, AVDA. ASTROFÍSICO FRANCISCO SÁNCHEZ, s/n, 38271 LA LAGUNA (STA. CRUZ DE TENERIFE), SPAIN
E-mail address: jbetanco@ull.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA, CAMPUS DE ANCHIETA, AVDA. ASTROFÍSICO FRANCISCO SÁNCHEZ, s/n, 38271 LA LAGUNA (STA. CRUZ DE TENERIFE), SPAIN
E-mail address: alecas000@gmail.com

INSTITUTO DE CIENCIAS MATEMÁTICAS, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, SERRANO 121, 28006 MADRID, SPAIN
E-mail address: jezabel.curbelo@icmat.es